

# Historical Background and Clinical Implications of Silver Diamine Fluoride in Dentistry: A Narrative Review

Neeraj Kumar<sup>1,\*</sup> and Richa Kumari<sup>2</sup>

<sup>1</sup>All India Institute of Medical Sciences, Bilaspur, Himachal Pradesh, India

<sup>2</sup>Ex-Tutor, ESIC Rohini, New Delhi, India

\*Corresponding author: [dr.neeraj001@gmail.com](mailto:dr.neeraj001@gmail.com)

**Abstract.** Dental caries is a chronic, irreversible disease affecting preschool children worldwide. As the frequency of dental caries has steadily declined over the past few years, children from low socioeconomic backgrounds still have higher rates of dental caries and still have a higher prevalence in developing countries. Though the role of fluoride in dental caries prevention has been scientifically proven, the antimicrobial effect of silver compounds to treat various infections has been postulated. However, 38% silver diamine fluoride (SDF) solution is currently used to arrest dental caries and as an antihypersensitivity agent. The SDF solution is available in dark, amber-colored bottles and is simple, noninvasive, and easy to use by trained dentists or specialists. Although it has the disadvantage of causing black discoloration and an unpleasant metallic taste after application, studies in the literature show that it can help prevent dental caries, hypersensitivity, root caries, and remineralization. This article explains in depth the SDF: its historical background, clinical application, and efficacy in different fields of dentistry.

**Keywords:** Remineralizing agent, silver diamine fluoride, caries prevention, topical agent, hypersensitivity.

## INTRODUCTION

Dental caries is a disease process that is multifactorial, dynamic, and irreversible that occurs due to an imbalance between host and ecological factors [1]. The epidemiological studies confirm that early childhood caries remains a significant disease burden in both developed and developing countries, along with high unmet dental needs [2]. The reasons for the unmet dental needs could be poor access to resources, financial constraints, the higher cost of dental treatment, lack of awareness, and their general health and education opportunities [3]. The traditional treatment for a carious tooth is to remove the infected, irreversible, and demineralized dentin by mechanical excavation and restore it with a restorative material [4]. In young children, behavior issues routinely complicate or prevent the restorative treatment. This unmet treatment needs lead to progression of the carious lesion and pain in some cases, which has a fatalistic impact on quality of life. Arresting the carious lesion has been proposed in children with uncooperative behavior and disadvantaged low socioeconomic communities [5].

Silver diamine fluoride (SDF), a colorless liquid with a pH of 8 to 9 and the ability to remineralize minerals, is utilized as a caries arresting agent in dentinal caries lesions in children. In the late 1970s, SDF was initially used in Japan but was not that popular. However, in the beginning of the 21st century, it has gained popularity in China as a dental caries arresting agent [6, 7]. It has recently become popular and available in the United States and South Asian countries. Other silver products like silver fluoride and SDF in different concentrations are commercially available in Austria, Argentina, and Brazil these days [8, 9] SDF is most commonly used in a concentration of 38% solution containing 44,800 parts per million (ppm) of fluoride. It is also available in 30% containing 35,400 ppm and 12% containing 14,150 ppm of fluoride, respectively [10]. The caries arresting efficacy of dental caries was reported to be 65.9% by 38% SDF [11]. The primary drawback of SDF is the black discoloration of dentinal carious tissue following application due to the oxidation process of silver ions, along with a few other reversible modifications such as oral tissue staining and ulceration [12]. This page provides information on this intriguing substance, including how it

can be used to treat and prevent dental problems in both primary and permanent teeth.

## HISTORICAL BACKGROUND


Although silver compounds were used to prevent dental caries, in Japan, it was customary for women to “Ohaguro,” or color their teeth black, to signify their marital status 1000 years ago [13]. Around 1000 B.C., the first medicinal use of silver compounds was to treat various infections. In medical science, silver-based materials such as silver foil, silver nitrate, and silver sutures have been employed [14]. Von Hofmann discovered silver nitrate to be a highly powerful antibacterial agent (Naegeli, 1893) [15]. In 1891, Stebbins used silver amalgam and nitric acid for the first time in dentistry and found caries inhibition in the subsequent follow-ups [16]. “Howe’s solution” consisting of silver nitrate was used directly on caries lesions and similar results were found by Howe et al. (1917) [17]. Since then, it has been used for caries arresting solution.

## SILVER DIAMINE FLUORIDE IN DENTISTRY

The first time silver fluoride was applied in dentistry was in the early 1970s [18]. Stannous fluoride ( $\text{SnF}_2$ ) must be used as a reducing agent for applying silver fluoride, a colorless, alkaline liquid that contains active silver and fluoride ions. SDF has been approved as a therapeutic agent by the Central Pharmaceutical Council of the Ministry of Health and Welfare, Japan dentistry [19]. SDF is a stable chemical compound, not as alkaline as silver fluoride (at a pH of 8–9) and does not require a reducing agent like silver fluoride. Practitioners in Southern California used SDF in preschool children with nursing bottle caries [20]. Many community projects were planned in sub-Saharan Africa, Cuba, and several African countries using SDF as a caries-arresting agent [5]. During this time, SDF is used quite seldom, and there has been limited research regarding its effectiveness. SDF has been used to stop tooth cavities in various nations, including China, Japan, the United States, and South Asian nations, at a concentration of 38%.

### Mechanism of Action

According to research (Lansdown, 2002a, 2006), silver interacts with sulphhydryl groups on proteins or DNA to block respiratory processes, change hydrogen bonding, affect the formation of cell walls, unwind DNA, and affect cell division at the microcellular level [21, 22]. Bactericidal activity and biofilm formation inhibition are produced by these interactions [23]. Shimizu and Kawagoe outline three potential modes of action (1976) [24]. Figure 1 illustrates the many therapeutic consequences of SDF.



**Figure 1.** Mechanism of action and clinical implications of SDF in dentistry.

### Obturation of dental tubules

Gottlieb (1947) hypothesized that dental caries can be prevented by obturation/blockage of dental tubules, i.e., blocking the organic invasive pathway [25]. When dentin is treated with SDF, silver and its derivatives stop acid diffusion and microbial penetration through the dental tubules. Shimizu (1974) showed that the dental tubules were shown to have reduced permeability and increased electric resistance. He further affirmed that the dentin’s surface area will shrink, and the dentin’s most readily demineralized region, the peritubular zone, may become coated with silver particles in obturated dental tubules [26]. In a recent study, Mei et al. (2013) discovered that the usage of 38% SDF protected the collagen from deterioration in demineralized dentin. The oligodynamic effect of the silver present in the dental tubules will restrict bacterial development and prevent demineralization of the dentin [27].

### Cariostatic action of the reaction products

Shimooka (1972) discovered that SDF ( $\text{Ag}(\text{NH}_3)_2\text{F}$ ) prevents and hardens dental cavities by reacting with hydroxyapatite crystals (HA) ( $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$ ) found in tooth material to produce calcium fluoride ( $\text{CaF}_2$ ) and insoluble silver phosphate ( $\text{Ag}_3\text{PO}_4$ ) [28].  $\text{CaF}_2$  continues to precipitate and hardens dental cavities by interacting with the insoluble silver phosphate crystals ( $\text{Ag}_3\text{PO}_4$ ) present in tooth material [28].

### Antibacterial properties through antienzymatic action

Suzuki (1976) suggested that the antibacterial properties of SDF are due to *Streptococcus mutans* cariogenic strains’

enzyme activity suppression and dextran-induced agglutination. We now provide two examples in this regard [32]. The reaction between silver and the organic component of dentin by ionophoresis leads to increased resistance to trypsin [33]. Additionally, Yanagida et al. (1971) demonstrated that collagenase and trypsin resistance in dentin protein treated with  $\text{Ag}(\text{NH}_3)_2\text{F}$  had risen [34].

## CLINICAL APPLICATION OF SILVER DIAMINE FLUORIDE

### Dental Caries Arresting Agent in Primary Teeth of Preschool Children

Most of the children in this study are suffering from severe early childhood caries. The treatment of these young children involves many difficulties, like child uncooperative behavior, so the majority of them remain left untreated. Nishino et al. (1969) and Moritani et al. (1970) compared two groups receiving SDF and without SDF and reported less caries increment in the children treated with SDF [35, 36]. In another study, SDF was used in a community-based caries control program, and results have shown that SDF effectively arrested the carious lesion in the primary teeth of preschool children [6]. In a systematic review, controlled clinical trials of caries prevention in primary teeth were evaluated by Oliveira BH et al. (2019) and concluded that SDF was effective compared to the fluoride varnish group or the placebo group in avoiding dental cavities [37]. According to a subsequent systematic study by Contreras (2017) [38], using SDF at concentrations of 30% and 38% is more successful in stopping dentinal caries in both primary and permanent dentition [38].

### Prevention of Pit and Fissure Caries

Due to structural differences, pits and fissures are more likely to develop dental cavities and are more difficult to keep clean than smooth surfaces. Pit and fissure caries are challenging to detect, and topical fluoride treatment has been shown to be less efficient in preventing pit and fissure caries. Due to its antibacterial properties, SDF has been demonstrated to be useful in reducing pit and fissure caries of the first molar teeth [39]. In a research by Nishino and Massler (1972), it was shown that the mean caries score was considerably lower in the fissures treated with silver fluoride or  $\text{SnF}_2$  compared to 8% in the SDF-treated group [40].

### Prevention of Secondary Caries

Most of the restorative materials are insoluble in oral fluids and adhere to the tooth structure. There are microspaces at the junction of the cavity wall and the restorative material, so microorganisms penetrate through microleakage, which leads to secondary caries [41]. SDF has been found to be

used for the prevention of secondary caries. Shimizu and Kawagoe (1976) in their study found that there was no evidence of secondary caries after 26 months of follow-up in primary teeth treated with SDF [24].

### Arrest Root Caries

Studies have shown a high prevalence of root caries in the elderly population [42–44]. The various studies mentioned that application of SDF arrests the root caries and there was less increment in the new carious lesions with the addition of SDF and oral health education [45, 46].

### Desensitize Sensitive Teeth

Gottlieb (1947) suggested that reaction products of SDF occlude the dentinal tubules, thus preventing dentin hypersensitivity and arresting dental caries [25]. Various studies have shown that SDF is more effective against chemical and mechanical wear such as abrasion, erosion, and hypersensitive dentin to thermal stimulation [47–49].

### Irrigation of Infected Root Canals

Silver compounds like ammoniated silver nitrate solution have been used as an irrigating solution to disinfect root canals. Silver fluoride solution has disinfectant and protein-coagulating properties. This blocks the root canal's dentinal tubules [50]. According to Hiraishi et al. (2010), 3.8% SDF is an effective antimicrobial root canal irrigant which removes the microbes from the canals and circum-pulpal dentin [51].

### Other Applications in Dentistry

The remineralizing action of SDF has been proven in the literature because of the fluoride content. When SDF, calcium sodium phosphate (CSP), and casein phosphopeptide-amorphous calcium phosphate complexes (CPP-ACP) were compared in an in vitro study by Vinod et al. (2020) using Diagnodent readings, the results revealed a statistically significant difference between baseline, post-demineralization, and post-remineralization readings. SDF demonstrated the highest levels of remineralization in the intergroup comparison, followed by CSP and CPP-ACP, respectively [52]. SDF also has antifungal potency against oral *Candida* species along with antimicrobial activity in the management of severe early childhood caries [53].

## DISADVANTAGES OF SILVER DIAMINE FLUORIDE

The main disadvantage and concern of SDF are black discoloration and metallic taste, so some parents and their children may not accept the treatment outcome. When SDF was applied to the carious dentin, insoluble yellow-colored

silver phosphate was formed [54], which later turns to black when exposed to sunlight or due to the influence of reducing agents. Knight et al. (2005) suggested application of after using SDF, potassium iodide is used to remove the black discoloration. Crystals of creamy white silver iodide are created when potassium iodide combines with the remaining free silver ions. As a result, there are no longer any free silver ions that may interact with sulfur to produce black precipitates [55].

SDF also has a metallic taste, which is unacceptable to the children and parents. Sometimes gingival and mucosal irritation can occur, which is transient in nature and the tissue affected turns white and heals within 1–2 days. Thus, use of a rubber dam or a separating medium like vaseline or cocoa butter should be used before using the SDF solution near to the gingiva [56].

Crystal et al. (2017) evaluated the parental acceptance and perception of SDF staining and observed that staining was more acceptable on posterior teeth compared to anterior teeth. Most parents preferred SDF application compared to treatment under deep sedation or general anesthesia [57]. In a different research, Alshammari et al. (2019) discovered that the majority of the parents disapproved of the SDF application. The authors came to the conclusion that dentists should offer informed consent forms with unambiguous photos while treating anterior teeth showing expected staining [58].

## CONCLUSION

SDF has been proven to be an effective therapeutic agent in arresting dental caries in both primary and permanent teeth. It also helps in reducing dentinal hypersensitivity, remineralizing properties, antimicrobial activity, and anti-fungal potency. Furthermore, SDF is a simple, easy to use, easily affordable, and widely available therapeutic agent in developing countries. Though it has the disadvantage of black staining and metallic taste, more studies would be needed to prove SDF as a material of choice in the current scenario.

## AUTHOR CONTRIBUTIONS

Both the authors contributed to concept designing, searching the literature, and writing the review article.

## ACKNOWLEDGMENT

The authors want to acknowledge that the article was solely written by the authors.

## ABBREVIATIONS

SDF, Silver diamine fluoride;  
SnF<sub>2</sub>, Stannous fluoride;

DNA, Deoxyribonucleic acid;  
ppm, Parts per million;  
HA, Hydroxyapatite crystals;  
CaF<sub>2</sub>, Calcium fluoride;  
CSP, Calcium sodium phosphosilicate; and  
CCP-ACP, Casein phosphopeptide-Amorphous calcium phosphate complexes

## REFERENCES

- [1] Fejerskov O., Kidd EAM., Nyvad B et al. (2008). Defining the disease: an introduction. In: Fejerskov O, Kidd E, editors. *Dental caries: the disease and its clinical management*. 2nd edition. Oxford (UK): BlackwellMunksgaard p. 4–6.
- [2] Edelstein BL. (2005). Pediatric caries Worldwide: implications for oral hygiene products. *References Compend Contin Educ Dent* 26, 4–9.
- [3] Baelum V., van PalensteinHelderma W., Hugoson A et al. (2007). A global perspective on changes in the burden of caries and periodontitis: implications for dentistry. *J Oral Rehabil* 34, 872–906.
- [4] Roberson TM., Heymann HO., Swift EJ. (2002). *Sturdevant's Art and Science of Operative Dentistry*. Mosby: St. Louis.
- [5] Bedi R. (1999). *Sardo-Infirri J. Oral health care in disadvantaged communities*, London: FDI world press.
- [6] Chu CH., Lo EC., Lin HC. (2002). Effectiveness of silver diamine fluoride and sodium fluoride varnish in arresting dentin caries in Chinese preschool children. *J Dent Res* 81, 767–770.
- [7] Llodra JC., Rodriguez A., Ferrer B et al. (2005) Efficacy of silver diamine fluoride for caries reduction in primary teeth and first permanent molars of school children: 36-month clinical trial. *J Dent Res* 84, 721–724.
- [8] Gotjamanos T. (1996). Pulp response in primary teeth with deep residual caries treated with silver fluoride and glass ionomer cement ('atraumatic' technique). *Aust Dent J* 41(5), 328–334.
- [9] Mei ML., Chin-Man Lo E., Chu CH. (2016). Clinical use of silver diamine fluoride in dental treatment. *Compend Contin Educ Dent* 37(2), 93–98.
- [10] Mei ML., Chu CH., Lo EC., Samaranayake LP. (2013). Fluoride and silver concentrations of silver diamine fluoride solutions for dental use. *Int J Paediatr Dent.* 23, 279–85.
- [11] Gao SS., Zhang S., Mei ML., Lo EC., Chu CH. (2016). Caries remineralisation and arresting effect in children by professionally applied fluoride treatment – a systematic review. *BMC Oral Health.* 16:12.
- [12] Rosenblatt A., Stamford TC., Niederman R. (2009). Silver diamine fluoride: a caries "silver-fluoride bullet". *J Dent Res.* 88, 116–25.
- [13] Yamaga R., Nishino M., Yoshida S et al. (1972). Diamine silver fluoride and its clinical application. *J Osaka Univ Dent Sch* 12, 1–20.
- [14] Russell AD., Hugo WB. (1994). Antimicrobiological activity and action of silver. *Prog Med Chem* 31, 351–370.
- [15] Von Naegeli V. (1893). *Dtsch Sci Schweiz NaturforschGes* 33, 174–182.
- [16] Stebbins EA. (1891). What value has argentinitras as a therapeutic agent in dentistry? *Int Dent J* 12, 661–670.
- [17] Howe PR. (1917). A method of sterilizing and at the same time impregnating with a metal affected dentinal tissue. *Dent Cosmos* 59, 891–904.
- [18] Craig GG., Powell KR., Cooper MH. (1981). Caries progression in primary molars: 24-month results from a minimal treatment programme. *Community Dent Oral Epidemiol* 9, 260–265.
- [19] Yamaga R., Yokomizo I. (1969). Arrestment of caries of deciduous teeth with diamine silver fluoride. *Dental Outlook* 33, 1007–1013.
- [20] Duperon DF. (1995). Early childhood caries: a continuing dilemma. *J Calif Dent Assoc* 23, 15–25.
- [21] Lansdown AB. (2002a). Silver I: Its antibacterial properties and mechanism of action. *J Wound Care* 11, 125–130.



- [22] Lansdown AB. (2006). Silver in health care: antimicrobial effects and safety in use. *CurrProbl Dermatol* 33, 17–34.
- [23] Wu MY., Suryanarayanan K., van Ooij WJ., Oerther DB. (2007). Using microbial genomics to evaluate the effectiveness of silver to prevent biofilm formation. *Water Sci Technol* 55, 413–419.
- [24] Shimizu A., Kawagoe M. (1976). A clinical study of diamine silver fluoride on recurrent caries. *J Osaka Univ Dent Sch* 16, 103–109.
- [25] Gottlieb B. (1947). *Dental Caries*, Philadelphia: Lea &Febiger. pp. 222–236.
- [26] Shimizu A. (1974). Effect of diammine silver fluoride on recurrent caries. *Jap. J. Conserv. Dent.* 17, 183–201. (in Japanese, summary in English)
- [27] Mei ML., Ito L., Cao Y., Li QL., Chu CH., Lo EC. (2013). The inhibitory effects of silver diamine fluorides on cysteine cathepsins. *J Dent.* 42(3), 329–335.
- [28] Shimooka S. (1972). On the penetration of silver nitrate and ammoniacal silver fluoride into microstructure of the sound dentin. *J. Nippon Dent. Coll* 59, 534–566. (in Japanese, summary in English)
- [29] Suzuki T., Nishida M., Sobue S et al. (1974). Effects of diamine silver fluoride on tooth enamel. *J Osaka Univ Dent Sch* 14, 61–72.
- [30] Kani T. (1970). X-ray diffraction studies on effect of fluoride on restoration of lattice imperfections of apatite crystals. *J. Osaka Univ. Dent. Soc* 15, 42–56. (in Japanese, summary in English)
- [31] Tamaki S. (1967). Effects of fluoride on crystallinity of synthetic and biological apatite. *J. Osaka Univ. Dent. Soc.* 12, 95–110. (in Japanese)
- [32] Suzuki T., Sobue S., Suginaka H. (1976). Mechanism of antiplaque action of diamine silver fluoride. *J Osaka Univ Dent Sch* 16, 87–95.
- [33] Sunada I., Kuriyama S., Komamura T et al. (1962). Resistance to acid and enzyme of dentin treated by metal ion ionophoresis. *Jap. J. Conserv. Dent* 5, 6–10. (in Japanese)
- [34] Yanagida I., Nishino M., Hano T et al. (1971). Effects of diammine silver fluoride on organic components of dentin of deciduous teeth. *Jap. J. Pedo.* 9, 39–46. (in Japanese)
- [35] Nishino M., Yoshida S., Sobue S et al. (1969). Effect of topically applied ammonical silver fluoride on dental caries in children. *J Osaka Univ Dent Sch* 9, 149–155.
- [36] Moritani Y., Doi M., Yao K et al. (1970). Clinical evaluation of silver diamine fluoride (Saforide) in controlling caries of deciduous teeth. *RinshoShika* 266, 48–53. (in Japanese)
- [37] Oliveira BH., Rajendra A., Veitz-Keenan A., Niederman R. (2019). The Effect of Silver Diamine Fluoride in Preventing Caries in the Primary Dentition: A Systematic Review and Meta-analysis. *Caries Res.* 53(1), 24–32.
- [38] Contreras V., Toro MJ., Elías-Boneta AR., AngelizEncarnación-Burgos. (2017). Effectiveness of silver diamine fluoride in caries prevention and arrest: a systematic literature review. *Gen Dent.* 65(3), 22–29.
- [39] Sato R, Sailo Y. (1970). Clinical Application of Silver Ammonia Fluoride (Saforide) to Children. *The Nippon Dental Review* 332, 66–67.
- [40] Nishino M., Massler M. (1972). Immunization of Caries-susceptible Pits and Fissures with a Diammine Silver Fluoride Solution, *The Japanese Journal of Pedodontics* 10, 6–11.
- [41] Anusavice K J. (2003). *Science of Dental Materials*, 11 th ed. Saunders Co., Philadelphia.
- [42] Chalmers JM., Hodge C., Fuss JM et al. (2002). The prevalence and experience of oral diseases in Adelaide nursing home residents. *Aust Dent J* 47, 123–130.
- [43] Lo EC., Luo Y., Dyson JE. (2004). Oral health status of institutionalised elderly in Hong Kong. *Community Dent Health* 21, 224–226.
- [44] Simunkovic SK., Boras VV., Panduric J et al. (2005). Oral health among institutionalised elderly in Zagreb, Croatia. *Gerodontolog* 22, 238–241.
- [45] Tan HP., Lo EC., Dyson JE et al. (2010). A randomized trial on root caries prevention in elders. *J Dent Res.* 89, 1086–1090.
- [46] Zhang W., McGrath C., Lo EC., Li JY. (2013). Silver diamine fluoride and education to prevent and arrest root caries among community dwelling elders. *Caries Res.* 47(4), 284–290.
- [47] Hatsuyama M., Toda T., Sckine I et al. (1967). Effect of Silver Ammonia Fluoride on Hypersensitive Dentin, *77re Journal of the Osaka Odontohgical Society* 1262–1265.
- [48] Murnse M., Takai H. (1969) Clinical Effect of Diamine Silver Fluoride on Cervical Hypersensitivity, *The Nippon Dental Review*; 323, 1123–1126.
- [49] Kimura K., Iso Y., Ohno. M et al. (1971). Clinical Test of Diammine Silver Fluoride (Saforide) Applied to Hypersensitive Dentin. *Shikagakuho* 71, 708–713.
- [50] Tanaka M. (1970). The effect of Silver Fluoride Applied in Pulp-removed Root Canal, *J. Osaka Univ. Dent. Soc.* 15, 34–47.
- [51] Hiraishi N., Yiu CK., King NM et al. (2010). Antimicrobial efficacy of 3.8% silver diamine fluoride and its effect on root dentin. *J Endod.* 36, 1026–1029.
- [52] Vinod D., Gopalakrishnan A., Subramani SM., Balachandran M., Manoharan V., Joy A. (2020) A Comparative Evaluation of Remineralizing Potential of Three Commercially Available Remineralizing Agents: An *In Vitro* Study. *International Journal of Clinical Pediatric Dentistry* 10.5005/jp-journals-10005-1715.
- [53] Fakhruddin KS., Egusa H., Ngo HC., Panduwawala C., Pesee S., Venkatachalam T., Samaranayake LP. (2020) Silver diamine fluoride (SDF) used in childhood caries management has potent antifungal activity against oral *Candida* species. *BMC Microbiology* 20(95), 2–11.
- [54] Halsted WS. (1895). The operative treatment of hernia. *Am J Med Sci* 110, 13–17.
- [55] Knight GM., McIntyre JM., Craig GG et al. (2005). An in vitro model to measure the effect of a silver fluoride and potassium iodide treatment on the permeability of demineralized dentine to *Streptococcus mutans*. *Aust Dent J* 50, 242–245.
- [56] Yamaga R., Nishino M., Yoshida S et al. (1972). Diamine silver fluoride and its clinical application. *J Osaka Univ Dent Sch* 12, 1–20.
- [57] Crystal YO., Janal MN., Hamilton DS., Niederman R. (2017) Parental perceptions and acceptance of silver diamine fluoride staining. *J Am Dent Assoc.* 148(7), 510–518.
- [58] Alshammari AF., Almuqrin AA., Aldakhil AM., et al. (2019). Parental perceptions and acceptance of silver diamine fluoride treatment in Kingdom of Saudi Arabia. *Int J of Health Sciences* 13(2).