

## Diagnosis and Protection of COVID-19 in Stomatology

Minzhi Yang<sup>1</sup>, Yujiong Chen<sup>1</sup>, Xiaoyan Hu, Mingsong Wu<sup>2,\*</sup> and Jianguo Liu<sup>2</sup>

<sup>1</sup>*School of Stomatology, Zunyi Medical University, Zunyi, Guizhou, China*

<sup>2</sup>*Special Key Laboratory of Oral Disease Research of Higher Education Institution of Guizhou Province, Zunyi Medical University, Zunyi, Guizhou, China*

\*Corresponding author: [mshwu0909@zmu.edu.cn](mailto:mshwu0909@zmu.edu.cn)

**Abstract.** In late 2019, a novel coronavirus uncertain etiology caused in Wuhan, China, and quickly spread to other provinces and countries. Some researchers isolated the pathogen named as the 2019 novel coronavirus disease (2019-nCoV). The common symptoms of patients infected with SARS-CoV-2, such as sore throat, high fever, diarrhea, and dyspnea. Besides that, researchers also recorded oral symptoms including ulcers, herpes, and tongue pain. At the same time, dental practitioners as a high-risk infection occupation always contact close patients. It is very important to prevent infection. The review aims to look back at the progress of the Coronavirus disease 2019 (COVID-19) associated with stomatology, to be studied, diagnosed, and prevented from the disease from the oral aspect.

**Keywords:** COVID-19, Oral, Diagnostic, Aged.

### INTRODUCTION

In December 2019, found the pneumonia of unknown source in China, it has broken out in many places around the world. This is a new zoonosis, which can be isolated from human airway epithelial cells, named 2019-nCoV [1]. The virus is an enveloped RNA virus [2], which possesses high infectivity and can cause fatal respiratory diseases. Because the clinical characteristics of COVID-19 are not significant and highly infectious, early diagnosis and prevention become critical to control the rapid spread of epidemic the outbreak. The common screening symptoms of the disease include fever, cough, sore throat, dyspnea, diarrhea etc, [3]. Furthermore, studies have shown that the main host cell receptor of SARS-CoV-2 is angiotensin-converting enzyme 2 (ACE2) [4]. The damage to various organs is linked to the distribution of ACE2 receptors in human systems. Therefore, ACE2 receptor distributed cells may become the host cells of the virus, causing the inflammatory response to related organs and tissues. ACE2 higher expression in oral mucosa may indicate that oral mucosa is a potential target of COVID-19, which may lead to oral mucosa associated infection, including ulcer, salivary gland inflammation, and so on, so that oral medical staff can make an early diagnosis of the patients through the oral cavity, to achieve

early prevention. Moreover, it is inevitable to generate fog, as well as aerosol, which makes doctors and patients exposed to a high-risk environment in dental diagnosis and treatment, so protective treatment is very crucial. The current research on COVID-19 related to oral disease is less reported, and it is easy to be ignored. This review mainly summarizes the research progress of suspicious symptoms, protective measures, and detection schemes through literature search, which attracts the attention of dental practitioners.

### SYMPTOMS AND SIGNS OF COVID-19

ACE2 is the target receptor for coronavirus to enter host cells, which are potential targets of SARS-CoV-2. At present, many studies reported some oral symptoms associated with COVID-19, including dry mouth, strange taste, oral mucosal lesions, and salivary gland inflammation. ACE2 may exists in oral tissues [5]. But, the correlation between new crown patients and oral symptoms is controversial.

#### Taste Dysfunction

Taste dysfunction is a common symptom of inpatients and most outpatients. Although other viral infections also

have altered taste functions, the incidence rate is considerably lower than that of COVID-19. It is reported that 88.8% of taste disorders (308 of 342 cases) in patients with a positive diagnosis of SARS-CoV-2 [6]. In a short case report, a couple was diagnosed as SARS-CoV-2 shortly after developing taste and olfactory dysfunction [7]. Moreover, 20.3% had the symptoms of taste or olfactory disorders before admission, among 88 patients diagnosed with COVID-19. The incidence of this symptom in women was higher than that in men [8]. What's more, the first symptom of covid-19 may be azoospermia, smell, and taste disorders [9]. There are currently two hypotheses about potential mechanisms that are reasonable: Firstly, the virus directly attacks olfactory neurons and damages its epithelium [10]. Secondly, the expression of ACE2 receptor was higher in the tongue than in another oral mucosa [5]. Lately, SARS-CoV-2 infection is expanding all over the world, which is a challenge to the medical system and medical resources. Sometimes, taste and smell can also be included in a screening condition. If these symptoms appear, patients can consider immediate home isolation.

### Oral Mucosal Lesions

Recently, few authors have reported that SARS-CoV-2 infected patients with common oral symptoms and signs, but the expression of ACE2 in the oral mucosa indicates the oral cavity may be a high risk site of COVID-19. A case report described an elderly male infected with COVID-19, who complained of pain from the palate, sore throat, and sore tongue, and other systemic symptoms such as fever, cough, and diarrhea. A complete intraoral examination found a large number of ulcers and blisters, resembled herpetic recurrent lesions, and subsequently diagnosed with COVID-19 [11]. Likewise, the other case reported 24 days after hospitalization, the internal examination found white plaque and several yellowish ulcers on the tongue dorsum. But when the patient's systemic symptoms improved, a few days after leaving the ICU, the oral symptoms quickly recovered [12]. Moreover, a study discovered the oral symptoms and signs, including ulcers that resembled simplex herpes lesion on their lip and tongue dorsum, denied that they had a history of recurrent ulcers disease, among 8 patients diagnosed with SARS-CoV-2 [13].

Although no oral tissue samples were obtained from patients with infected COVID-19 in the actual study, the author found that the evolution of oral lesions paralleled the clinical symptoms of COVID-19 [13]. It is very vital to know whether SARS-CoV-2 destroys oral mucosa epithelium, then causes oral ulcer or plaque, and what is a new pathogenic mechanism. What's more, we could assume whether oral lesions are the first symptom caused by COVID-19 infection.

### Sialadenitis

A reported that the expression of ACE2 was higher in salivary glands epithelial cells than in lung cells, explaining that salivary glands also may be a potential target of COVID-19 [14]. Therefore, we should not ignore the potential infection from saliva. The virus damaged the gland, leading to salivary gland inflammation, including submandibular gland inflammation and parotitis. Furthermore, it was reported that SARS-CoV-2 could be detected in saliva before in lung lesions [15]. In conclusion, the virus of patients with asymptomatic carriers may originate from saliva generated by the salivary gland. Many researchers have hypothesized that salivary glands infection may change in salivary flow rate and composition, which cause taste alterations and dry mouth [16].

### ORAL MANIFESTATIONS IN THE ELDERLY WITH UNDERLYING DISEASES

Mueller et al. found that hospitalized patients infected with SARS-CoV-2 were over 65 years old accounted for 80%, and the mortality of elderly patients is also higher than that of young people [17]. Some studies show that many basic diseases in elderly patients with cardiovascular diseases, hypertension, diabetes, and chronic obstructive pulmonary disease (COPD) are the major risk factors for the increase of COVID-19 and main life-threatening risk factors [18]. The cytokines and microorganisms released by oral infection can cause distal organ inflammation, so oral health is associated with general health. And local oral disease may promote the severity of COVID-19 in older. A recent study, hypertension, obesity, and diabetes were identified as the three most potential adverse outcomes in COVID-19 patients requiring hospitalization [19]. The elderly obese patients are prone to oral diseases, especially gingivitis and periodontitis because obesity will change the composition of periodontal microorganisms and increase pathogens. Obesity can reduce immunity, cause serious damage to organs and tissues, and cause complications of a variety of systemic diseases [20]. Compared with normal weight, adipose tissue of obese patients secretes pro-inflammatory factors and adipokines, which will aggravate the inflammatory effect of periodontitis and SARS-CoV-2 infection [21]. Studies have found that obese people are seemly to be possible to be infected with COVID-19, among which co-diseased patients with periodontitis and obesity may increase the risk of infection in obese patients [22]. In addition, it has been reported that the level of chronic inflammatory marker C-reactive protein is increased not only in hospitalized patients with COVID-19, but also expressed in patients with periodontitis, even in obese patients with periodontitis. With the increase of age, the prevalence of obesity and periodontal disease will

increase, which increases the prevalence and death risk of new crowns.

It also reported that 12% of the 140 patients with COVID-19 were diabetic. Scholars believe that the elderly patients with diabetics are more likely to infect the virus, even as a predictor of morbidity and mortality of periodontal disease and COVID-19 [23]. Studies have found the expression of ACE2 is significantly increased in diabetic patients, on the one hand because patients with diabetes and hypertension will increase angiotensin II type I receptor blockers (ARBs) and ACE inhibitors [24]. Moreover, COVID-19 may cause inflammation through ACE2 imbalance and cause complications of diabetes. Meanwhile, virus entering cells can delay death, evade host immunity, and increase the replication and secretion of chemokines. When infected with SARS-CoV-2, immune response and interferon will decrease, proinflammatory factors will increase, causing cytokine storm, leading to serious local tissue damage, such as periodontitis.

Hypertension is a common senile disease and one of the comorbidities of infection with SARS-CoV-2. It has been reported that comorbid hypertension is the highest risk factor in 1099 hospitalized patients with new crown infection [24]. ACE inhibitors and angiotensin receptor blockers (ARBs) are used to treat hypertensive patients, but both of which will increase ACE2 expression [25]. Therefore, increased ACE2 expression in hypertension and periodontitis may indicate an increased risk SARS-CoV-2 infection.

Cancer is also considered to be one of the risk factors for SARS-CoV-2 infection and aggravation because of its systemic immune suppression. When a systemic immune disorder is accompanied by the increase of cytokines and chemokines, it promotes inflammation and accelerates the serious progress of COVID-19, and leads to a local inflammatory response in the oral cavity, especially periodontitis. When the host is immunosuppressed, the interaction between virus and host epithelial cells will destroy the integrity of tissue and cause local lesions, such as necrotizing gingivitis, periodontitis, and dry mouth.

Oral diseases caused by systemic basic diseases of COVID-19 comorbidity, oral local inflammatory comorbidity covid-19 will also cause systemic lesions. For example, persistent periodontal disease will cause the release of inflammatory cytokines and enter the systemic circulation, induce systemic inflammation, and aggravate the viral infection of covid-19. There are many systemic diseases in the elderly. Due to the poor oral hygiene environment, the oral bacterial load increases, which destroys the symbiotic relationship between oral microorganisms. Bacteria can be colonized in distal organs through blood, and even spread to the respiratory tract through the inhalation of oral liquid during breathing, increasing the risk of pneumonia or respiratory diseases [26]. Other scholars hypothesized that gram-negative periodontal bacteria from the mouth may lead to lipopolysaccharide-induced

lung cell aging and accelerate SARS-CoV-2 replication [26].

Therefore, it is very crucial to pay attention to the oral health of the elderly, specifically the elderly in the intensive care unit. Improving oral health can reduce complications and COVID-19 incidence of distal organs. Although there is no clear causal relationship between oral cavity, systemic diseases, and COVID-19 in the elderly, oral-related inflammation can cause microbial ecological imbalance, host stress response, and immune imbalance to stimulate the severity of COVID-19.

## DEPARTMENT OF STOMATOLOGY PROTECTION DURING COVID-19

Due to the highly infectious, and worldwide epidemic, COVID-19 is a new challenge to the stomatology department. There are several characteristics in oral treatment such as long treatment time, closely contact between patients and doctors, the water mist and aerosol formed by saliva, blood, as well as secretion of patients during the operation of dental instruments. The aerosol can spread up to 2.1 meters, and splash into the face, eyes, or blood of medical staff or other patients [27]. Furthermore, the liquid, materials, and contaminated dental instruments contacted by dental professionals may also be the route of virus transmission. Sometimes, a dental clinic is one of the most concentrated places for patients. It is possible that patients conceal their illness or potential infectious diseases, which may cause a big outbreak. Therefore, protection has become very indispensable in the department of stomatology.

### Personal Protective Equipment Department of Stomatology

Personal protective equipment department of stomatology is a high exposure occupation during the novel coronavirus disease 2019, so occupation protection is very important. (1) Researchers investigated the protection status of medical staff in the department of stomatology through questionnaires. The results showed that 80.71% of the 1229 medical staff used ordinary masks. The wearing rate of N95 masks was only 0.56%. Only 67.61% of the medical staff changed masks in 4~6 hours. Respirator has an 85% chance of not being infected with the virus in high-risk condition [28]. In the past few months, Ma et al., analyzed the effect of N95 masks, family masks, and surgical masks on virus isolation, and showed that N95 masks were greater reliability [27]. (2) Gowns and gloves: In a dental setting with a severe infection rate, to increase the protection function, the professionals should use disposable protective clothing. Medical disposable protective clothing should cover their head, trunk to ankle, and protect all parts of the body. Furthermore, it should not be able to penetrate the virus-carrying saliva and blood when splashed on the

protective clothing during a dental operation. Professionals are prone to cross-infection in practice, so they must wear protective gloves and change them for each patient. Besides this, gloves should be replaced when contacting the cleaning area. (3) Hand hygiene: the first measure reduce cross-transmission of the virus in a dental environment [29]. Washing hands with soapy water and alcohol is particularly important to control the spread of respiratory diseases including SARS [30]. Due to the transmission characteristics of the 2019-nCoV virus, a study suggests the “two before and three after” technique as standard hand hygiene procedures. Doctors must wash hands before examining the patient, before dental practices, after contact with the patient, after touching the environment without previous disinfection, and after touching the oral mucosa and skin of a patient. Furthermore, WHO recommends that professionals should frequent hand-washing before and after any direct or indirect contact with the patient [31].

### Prevention and Control of Hospital

The dental clinic has always been a high-density place with a high flow of people, so it is very important to do a good job in the prevention and control of nosocomial infection during this time. Otherwise, it is easier to increase the risk of infection. About the prevention and control of nosocomial infection, we summarized the following points through literature inquiry. First, we advocated orderly consultation and diagnosis before treatment, including telephone or remote video consultation and diagnosis, to preliminarily screen whether patients can go to the outpatient clinic, and formulate a treatment plan and preventive nursing. It also includes emergency priority to reduce the number of outpatient visits. The department of stomatology in many countries advocates the principle of oral emergency priority. During the outbreak of this disease in early 2020, China only opened oral emergency to deal with some acute toothache, acute space infection, tooth trauma and so on. Likewise, British Medical Journal stipulates that ibuprofen will not be used to treat toothache, but paracetamol will be selected as a new analgesic to treat some simple toothache. The main reason is that ibuprofen can interfere with the function of the human immune system [32]. Furthermore, a survey before visiting time survey was conducted by the New Zealand Dental Association and other associations, included whether COVID-19 positive or respiratory symptoms were noted: cough, shortness of breath, sore throat, travel outpatient history and body temperature, make dentists understand whether or not COVID-19 might be exposed to an earthquake. If the temperature exceeds 37.5°C and is accompanied by respiratory disease-related symptoms, the dental visit will be delayed for 14 days [33]. At the time you make the appointment, to reduce the amount of oral microorganisms in saliva, the researchers suggest that 0.2% povidone-iodine or 1% hydrogen peroxide gargle be used before dental

practice [34]. In addition, since the bacterial aerosols are the largest at 1m from the headrest, it is found that rubber dam isolation can effectively reduce the contaminated gas produced by dental treatments [35]. In the meantime, WHO suggests that sodium hypochlorite should be used to disinfect the public areas and appliances in the dental clinic [31].

## DISCUSSION

Although many countries have already developed and started to use efficient COVID-19 vaccine, SARS-CoV-2 is spreading in global. Even in a short time, experts assess that the epidemic will continue. Early detection, diagnosis and isolation and treatment are necessary to outbreak control measures. We urgent need to seek for one safe, sensitive and highly efficient method to detect the rival. In addition, dental clinics have a high density of patients, have more transmission routes. Therefore, practicing dentists should not ignore any oral symptoms of the development and changes of patients tested positive for COVID-19 during the period. The review has some limitations. There is a limited number of primary reports and data about dental research related to COVID-19. Likewise, the studies of pathology and mechanism of oral symptoms of patients infected with COVID-19 are still unclear. What a pity that the vaccines of SARS-CoV-2 are entering various regions and countries, but few scholars have reported changes in oral protection after the entry of vaccines. In short, more scholars in the field of Stomatology are needed to fully research the oral symptoms, manifestations, and pathogenesis of COVID-19 infection in the future, and timely evaluate and treat them, to improve the quality of life of patients.

## CONFLICT OF INTEREST

No conflict of interest in this work.

## AUTHOR CONTRIBUTIONS

Minzhi Yang: Data curation and writing-original draft, including: preparation, creation and presentation of the published work, specifically writing the initial draft.

Yujiong Chen: Fromal analysis, application of statistical, mathematical, computational, or other formal techniques to analyze or synthesize study data.

Mingsong Wu: Writing-Original graft, preparation, creation and presentation of the published work by those from the original research group, specifically critical review, commentary or revision-including pre-or postpublication stages.

Jianguo Liu: Oversight and leadership responsibility for the research activity execution.

## FUNDING

This study was supported by the Construction Projects of Medical Biomaterial Research & Development Talent Base in Guizhou Province and Zunyi City (No. 3, and No. 69), and Science and Technology Project of Guizhou Province (5772-006).

## REFERENCES

- [1] Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med*, 2020; 382, 727–733.
- [2] van Regenmortel MH, Mayo MA, Fauquet CM, et al. Virus nomenclature: consensus versus chaos. *Arch Virol*, 2000; 145, 2227–2232.
- [3] Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *Jama*, 2020; 323, 1061–1069.
- [4] Wu F, Z7 hao S, Yu B, et al. A new coronavirus associated with human respiratory disease in China. *Nature*, 2020; 579, 265–269.
- [5] Xu H, Zhong L, Deng J, et al. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int J Oral Sci*, 2020; 12, 8.
- [6] Lechien JR, Chiesa-Estomba CM, De Siati DR, et al. Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch Otorhinolaryngol*, 2020; 277, 2251–2261.
- [7] Hjelmæsæth J, Skaare D. Loss of smell or taste as the only symptom of COVID-19. *Tidsskr Nor Laegeforen*, 2020; 140.
- [8] Giacomelli A, Pezzati L, Conti F, et al. Self-reported Olfactory and Taste Disorders in Patients With Severe Acute Respiratory Coronavirus 2 Infection: A Cross-sectional Study. *Clin Infect Dis*, 2020; 71, 889–890.
- [9] Paderno A, Schreiber A, Grammatica A, et al. Smell and taste alterations in COVID-19: a cross-sectional analysis of different cohorts. *Int Forum Allergy Rhinol*, 2020; 10, 955–962.
- [10] Baig AM, Khaleeq A, Ali U, et al. Evidence of the COVID-19 Virus Targeting the CNS: Tissue Distribution, Host-Virus Interaction, and Proposed Neurotropic Mechanisms. *ACS Chem Neurosci*, 2020; 11, 995–998.
- [11] Sinadinos A, Shelswell J. Oral ulceration and blistering in patients with COVID-19. *Evid Based Dent*, 2020; 21, 49.
- [12] Amorim Dos Santos J, Normando AGC, Carvalho da Silva RL, et al. Oral mucosal lesions in a COVID-19 patient: New signs or secondary manifestations? *Int J Infect Dis*, 2020; 97, 326–328.
- [13] Brandão TB, Gueiros LA, Melo TS, et al. Oral lesions in patients with SARS-CoV-2 infection: could the oral cavity be a target organ? *Oral Surg Oral Med Oral Pathol Oral Radiol*, 2021; 131, e45–e51.
- [14] Xu J, Li Y, Gan F, et al. Salivary Glands: Potential Reservoirs for COVID-19 Asymptomatic Infection. *J Dent Res*, 2020; 99, 989.
- [15] Liu L, Wei Q, Alvarez X, et al. Epithelial cells lining salivary gland ducts are early target cells of severe acute respiratory syndrome coronavirus infection in the upper respiratory tracts of rhesus macaques. *J Virol*, 2011; 85, 4025–4030.
- [16] da Silva Pedrosa M, Sipert CR, Nogueira FN. Altered taste in patients with COVID-19: The potential role of salivary glands. *Oral Dis*, 2021; 27 Suppl 3, 798–800.
- [17] Mueller AL, McNamara MS, Sinclair DA. Why does COVID-19 disproportionately affect older people? *Aging (Albany NY)*, 2020; 12, 9959–9981.
- [18] Wang B, Li R, Lu Z, et al. Does comorbidity increase the risk of patients with COVID-19: evidence from meta-analysis. *Aging (Albany NY)*, 2020; 12, 6049–6057.
- [19] Richardson S, Hirsch JS, Narasimhan M, et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *Jama*, 2020; 323, 2052–2059.
- [20] Palmer AK, Xu M, Zhu Y, et al. Targeting senescent cells alleviates obesity-induced metabolic dysfunction. *Aging Cell*, 2019; 18, e12950.
- [21] Peters U, Dixon AE, Forno E. Obesity and asthma. *J Allergy Clin Immunol*, 2018; 141, 1169–1179.
- [22] Larvin H, Wilmott S, Kang J, et al. Additive Effect of Periodontal Disease and Obesity on COVID-19 Outcomes. *J Dent Res*, 2021; 100, 1228–1235.
- [23] Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72,314 Cases From the Chinese Center for Disease Control and Prevention. *Jama*, 2020; 323, 1239–1242.
- [24] Li XC, Zhang J, Zhuo JL. The vasoprotective axes of the renin-angiotensin system: Physiological relevance and therapeutic implications in cardiovascular, hypertensive and kidney diseases. *Pharmacol Res*, 2017; 125, 21–38.
- [25] Ferrario CM, Jessup J, Chappell MC, et al. Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2. *Circulation*, 2005; 111, 2605–2610.
- [26] Aquino-Martinez R, Hernández-Vigueras S. Severe COVID-19 Lung Infection in Older People and Periodontitis. *J Clin Med*, 2021; 10, 279.
- [27] Ma QX, Shan H, Zhang HL, et al. Potential utilities of mask-wearing and instant hand hygiene for fighting SARS-CoV-2. *J Med Virol*, 2020; 92, 1567–1571.
- [28] MacIntyre CR, Chughtai AA, Rahman B, et al. The efficacy of medical masks and respirators against respiratory infection in healthcare workers. *Influenza Other Respir Viruses*, 2017; 11, 511–517.
- [29] Larson EL, Early E, Cloonan P, et al. An organizational climate intervention associated with increased handwashing and decreased nosocomial infections. *Behav Med*, 2000; 26, 14–22.
- [30] Fung IC, Cairncross S. Effectiveness of handwashing in preventing SARS: a review. *Trop Med Int Health*, 2006; 11, 1749–1758.
- [31] WHO Guidelines Approved by the Guidelines Review Committee. *Infection Prevention and Control of Epidemic- and Pandemic-Prone Acute Respiratory Infections in Health Care*. Geneva: World Health Organization Copyright © World Health Organization 2014; 2014.
- [32] Ather A, Patel B, Ruparel NB, et al. Coronavirus Disease 19 (COVID-19): Implications for Clinical Dental Care. *J Endod*, 2020; 46, 584–595.
- [33] Gurgel BCV, Borges SB, Borges REA, et al. COVID-19: Perspectives for the management of dental care and education. *J Appl Oral Sci*, 2020; 28, e20200358.
- [34] Peng X, Xu X, Li Y, et al. Transmission routes of 2019-nCoV and controls in dental practice. *Int J Oral Sci*, 2020; 12, 9.
- [35] COVID Samaranyake LP, Reid J, Evans D. The efficacy of rubber dam isolation in reducing atmospheric bacterial contamination. *ASDC J Dent Child*, 1989; 56, 442–444.