

Molecular Technique for Gender Identification: A Boon in Forensic Odontology

Rahul Pandey¹, Shruti Singh², Ruby Dwivedi³, Ekta Anthony⁴, Divya Mehrotra⁵
and Shaleen Chandra^{6,*}

¹Scientist, DHR-MRU, King George's Medical University, Lucknow, UP, India

²Senior Resident, Department of Oral Pathology and Microbiology, King George's Medical University, Lucknow, UP, India

³Women Scientist, DHR-MRU, King George's Medical University, Lucknow, UP, India

⁴PhD Scholar, DHR-MRU, King George's Medical University, Lucknow, UP, India

⁵Professor, Department of Oral and Maxillofacial Surgery, King George's Medical University, Lucknow, UP, India

⁶Professor and Head, Department of Oral Pathology and Microbiology, King George's Medical University, Lucknow, UP, India

*Corresponding author: profshaleenchandra@gmail.com

Abstract. Identification of an individual is essential both for legal and humanitarian purpose. Gender identification is an essential criteria required to establish identity of an individual. Use of molecular technique in forensic odontology is an emerging field and widens the avenue even in challenging cases of identification. Gender determination using molecular techniques in forensic odontology shows prodigious outcome and hence befitted as frontier in forensic odontology. Identification of an individual in mass disaster and crime investigation cases in current scenario is an ardent but highly desirable task. Thorough knowledge of forensic genetics is essential and is the need of the hour. Recent advancements in molecular technique for DNA typing useful in gender determination are both economical and time savers. These advancements will bring justice to people and large benefit to the society. An exhaustive internet based literature search was carried out in pubmed and google scholar database using terms 'gender identification' OR 'Individual identification' AND 'DNA fingerprinting'. The search yielded 1848 articles. Inclusion criteria included relevance to the context decided by reading the abstract and availability of full text. A total of 169 articles were selected. Further on reading the full texts, only article pertaining to oral cavity along with relevant articles from the references of the selected articles were considered. A total of 90 articles were considered in the final review. The current paper aims to comprehensively review the gender determination molecular techniques useful in forensic odontology. It emphasizes on various sources, techniques, profiling systems and genes employed for gender identification from oral cavity.

Keywords: Gender determination, Forensic odontology, Molecular techniques, Amelogenin, Alphoid repeats.

INTRODUCTION

Personal identification is prime requisite in several instances including mass disasters resulting from natural calamities (earthquakes and floods), terrorist attack or mass murders and other situations where the biological

remains are insufficient or decomposed enough to cause difficulty in identification [1].

Personal identification relies on features aiding in tentative identification described as 'big fours' (determination of age, **gender**, stature and ethnicity). This is required for legal, social and humanitarian purpose.

Gender determination forms the pillar of personal identification. The conventional method for gender determination includes analysis of morphological differences in the various anatomical parts such as bone, mandible, clavicle, skull etc. [2] Though the efficiency in individual or gender identification using morphological features is appreciable but sole dependence on the same is hampered in cases where complete tooth or desired structure is unavailable. In such cases, employing molecular techniques proves a boon. Decoding of individual DNA via DNA fingerprinting or DNA profiling aids in individual and gender identification.

This review aims to comprehensively review the gender determination molecular techniques useful in forensic odontology. It emphasizes on various sources, techniques, profiling systems and genes employed for gender identification from oral cavity.

METHOD OF DATA COLLECTION

An exhaustive internet based literature search was carried out in pubmed and google scholar database using terms 'gender identification' OR 'Individual identification' AND 'DNA fingerprinting'. The search yielded 1848 articles. Inclusion criteria included relevance to the context decided by reading the abstract and availability of full text. A total of 169 articles were selected. Further on reading the full texts, only article pertaining to oral cavity along with relevant articles from the references of the selected articles were considered. A total of 90 articles were considered in the final review.

MORPHOLOGICAL AND MOLECULAR BASIS FOR GENDER DETERMINATION

Gender determination plays a key role in personal identification. Several morphological characteristics exhibit sexual dimorphism thereby aiding in identification of the gender of the individual. Forensic dentists analyze key morphological differences in bones particularly skull including mandible depending on the reported literature findings. (Refer Table 1).

It is well known that tooth escapes most insults in adversities like fire, explosions and decomposition of bodies or situations with minimal forensic biological sample availability. Forensic odontology comes into play to support human identification in such scenarios. (Refer Table 2).

Gender identification via differentiating morphological features is quite efficacious but in cases with incomplete or destructed dental structures, amalgamation of Forensic odontology with molecular techniques comes to rescue. Decoding of individual DNA via DNA fingerprinting or DNA profiling aids in individual and gender identification [3].

Analysis of DNA is relatively easy to work with owing to its biologically robust molecular structure being capable

Table 1. Differentiating features in the skull for sexual dimorphism.

Characteristic Feature	Male	Female
Skull Architecture	Big and rugged	Sloped and smooth
Cranial mass	More blocky And massive	Rounder and tapers at the top
Temporal Ridge	More prominent	Less prominent
Supraorbital Margin	Round and dull	Sharper
Eye Orbit	Sharp superior margin	Circular, higher and larger
Frontal lobe	Low and sloped	Higher and rounded
Zygomatic Bone	More pronounced	Less pronounced
Super ciliary Arch	Larger and pronounced	Smaller
Gonion	Flared and sharply angled	Less flared
Teeth	Larger	Smaller
Mastoid	Medium to large	Small to medium
Nasal Aperture	High and sharp	Lower and round

of procurement and analysis even from archival specimens like hair, dried blood, fixed tissue sections or even exfoliated cells into secretions such as saliva.

Also the unique nucleotide sequences present in DNA of biological tissue result in differential genetic expression and variations in protein levels leading to physical and functional variations between individuals revealing an individual's DNA constitution and therefore reveal person's identity. The DNA profile tests are completely consistent and are authentic legal evidence required in paternity disagreements as well as identification of individuals [4-6].

ROLE OF DNA FINGERPRINTING IN IDENTIFICATION

After its discover by Watson and Crick in 1953 [4], the double-helix structured DNA has been enormously studied in almost all field of sciences since it provides the blueprint of the genetic make-up of every species. DNA has exons (protein coding segments constituting 2-5% of total cellular DNA) and introns (Non-coding DNA comprising the remaining 95%). The introns owing to no other known function apart from acting as a spacer DNA between exons are referred as junk DNA. It generally exists in multiple copies leading to repetitive DNA (20-30%) [5]. These regions are unique and highly polymorphic (different in each individual). According to their size, these repeats are labeled as variable number of tandem repeat (VNTR) also called as mini-satellites (9-80 base-pairs), and short tandem repeats (STR) popularly known as microsatellites (2 to 5 bp) [6]. In addition to this,

Table 2. Characteristic Dental features for sexual dimorphism.

	Characteristic	Male	Female
METRIC ANALYSIS	Mesio-distal and Bucco-lingual Dimension.	More	Less
	Canine dimorphism	More	Less
	– Inter-canine width		
	– Mesio-distal width		
	Dental Index	More	Less
	– Incisor index		
	– Canine Index (Mandibular Canine)		
NON METRIC	Distal Accessory ridge of canine	More pronounced	Less pronounced
	Number of cusps in mandibular first molar	4 cusp molar	5 cusp molar
	Shape of mandible	Square	Round
	Maxillary Sinus	Large volume and width	Small volume and width
RUGAE ANALYSIS	Rugae length	Short	Long
	Fragmentary Rugae	Less	More
LIP PRINTS		Type III, IV, V	Type I, I', II

single base variations in DNA sequence called polymorphisms also exist.

Alec Jeffrey (1985) described 'DNA fingerprinting' or DNA typing (profiling) for deciphering these variations for confirming an individuality [7]. DNA fingerprinting depends upon prompt recognition by probe along with constant Mendelian pattern of inheritance. DNA profiling is sensitive, reproducible, amenable to computer data basing, and therefore the forensic DNA technology is fore-runner in deciphering complex criminal cases and, also solving paternity disputes all over the world [8].

EXTRACTION OF DNA FROM ORAL CAVITY

Although whole blood is the highest yielding source of DNA, it may not be accessible in certain adverse circumstances. In such cases, DNA can be appreciably extracted from various sources in the oral cavity for example teeth, saliva or saliva stained samples, buccal swabs. Also DNA from cells attached to toothbrush, oral prosthesis such as dentures has been successfully obtained.

Lessing et al. (1995) demonstrated use of dental parameters and extracted DNA from dental tissues for individualization and later Malaver et al. (2003) studied samples from cadavers buried for more than five years and extracted DNA from pulp, dentin and cementum from the same. DNA can be obtainable from pulp of the teeth that had been kept in various environmental conditions [9, 10]. Appreciable amount of DNA can be isolated from teeth, buccal swabs, saliva and saliva stained samples. Isolation of DNA from buccal swabs is quite cost-effective method due to ease in sample collection, long-term storage capability and minimal amount of sample requirement to obtain sufficient DNA [11]. Ghatak et al established simplified phenol-chloroform method for DNA extraction using from buccal swabs [12].

Walsh et al. (1992) successfully extracted DNA from saliva for the first time. Since then several studies demonstrated saliva and salivary stained samples as suitable source of DNA. Saliva and salivary stained sample contain desquamated epithelial cells from the lips and oral mucosa and is relatively easy to collect, painless and non-invasive procedure [13, 14].

GENDER DETERMINATION VIA DNA FROM ORAL CAVITY

Highly degraded human remains present a great challenge in identification. Classical method of gender identification includes visual analysis of the bones having characteristic pattern for each gender. But, such techniques are unsuitable in situations where the samples are insufficient or extremely damaged. Therefore, alternate technology for sex determination with molecular biology has emerged as one of the best methods and hence has garnered immense attention.

As discussed earlier, DNA extraction from oral cavity has been the most common method employed for gender identification. Literature reveals integration of newer protocols and technologies that have led to better results for DNA extraction. Anzai-Kanto et al. recovered DNA followed by its typing from saliva that was deposited on skin of the volunteers in order to assess its potential for forthcoming case analysis [15].

Identification of gender employing SRY gene amplification using real-time PCR from isolated epithelial cells of removable partial denture has also been attempted and dentures stained with saliva proved to be excellent source for forensic DNA [16]. However, limitations with the saliva include small amount usually deposited on the skin particularly in bite impressions. Moreover, there are potential chances of cross contamination and mixing leading to aberrant results.

Apart from saliva buccal swabs could be another potential source of DNA. But limitations include unavailability of buccal swabs in people who died long before, potential

Table 3. Gender determination studies using molecular techniques from Oral Cavity samples.

Studied by	Study Samples, Design and Description	DNA Primer Used	Outcome
Yamada et al. (1989) [55]	Tooth samples (Dental Pulp)	Using 32 P labeled DNA probe 'Myo' and Y-chromosome specific DNA Probe	DNA extracted from dental pulp is useful for identification of individuals, paternity testing, and sex determination.
Potsch et al. (1992) [7]	Freshly extracted teeth stored for 4 years and 6 weeks at room temperature (n = 30); teeth from jaw fragments stored for 15 years; teeth extracted from actual identification cases (n = 8)	Genomic dot blot hybridization using biotinylated repeated DNA probe PHY2.1	Sex was accurately identified in all cases using 50–100 ng target DNA.
Chen et al. (1994) [63]	Human extracted teeth subjected to different conditions; varying pH solution; 10% formalin, 75% ethanol; different temperature (–20 degrees C, 4 degrees C, 25 degrees C, 37 degrees C, boiling, burning); humidity (20%, 66%, water submergence); burying the teeth outdoors; and aging (teeth stored at room temperature for 1.5, 3.5 and 8 years).	Gender determination using probe Y1.1 and Y 1.2	154 bp fragment was obtained in all male teeth, No 154 bp fragment was found in a female fresh tooth. 154 bp fragment bands were not observed in case of teeth subjected to burial outdoors for 10 weeks, submergence in water for 10 weeks, soak in pH 2 solution for 10 weeks, and combustion for 10 minutes; only faint bands were observed in case of teeth subjected to 66% humidity and pH 10 solution for 10 weeks and combustion for 5 minutes; obvious bands were observed in the rest cases. The quality of DNA from dental pulp affected PCR amplification result.
Pawlowski et al. (1995) [61]	Gender determination from Fomalin fixed paraffin embedded tissues (FFPE)	Amelogenin	It is possible to identify using 788 bp and 977 bp amelogenin fragment from soft tissues stored for 2 years.
Yomamoto et al. (1996) [62]	DNA profile typing from dental pulp	Y specific Probe	DNA typing useful for accurate sex determination from extracted teeth upto 21 months.
Hanaoka et al. (1996) [51]	Sex determination from blood and teeth	Alphoid repeat primers	Gender determination could be done with precision.
Pilly et al. (1997) [47]	Sex determination from pulp from 21 male and 24 female using third molar teeth	ZFX and ZFY	Males distinguished from female by 317 base pairs.
Jiang et al. (1997) [64]	Gender determination using DNA obtained from various samples (including saliva)	Amelogenin	Sensitive, rapid and reliable gender determination from various sample could detect gender in as little as 50 pg DNA.
Ouchinniko et al. (1998) [65]	Archeological samples from teeth samples without any cavities (n = 15)	Amelogenin (DYZ1, DYZ3, DXZ3 loci)	Successfully determined gender from all samples.
Komuro et al. (1998) [58]	Gender determination from dental pulp by capillary gel electrophoresis (n = 10 males and 10 females)	Amelogenin	Correct Gender determination from all cases.
Urbani et al. (1999) [66]	To determine effect of temperature on sex determination from DNA PCR analysis from human dental pulp (n = 94)	Amelogenin	PCR analysis was 100% reliable, Gender assessment from teeth heated at 100°C for 15 minutes but less reliable when the teeth were heated at higher temperatures for longer periods of time.
Meyer et al. (2000) [21]	Gender determination from three tooth samples (ancient)	Amelogenin	Success rate >90%.
Inoue et al. (2000) [67]	They determined gender from 55 resin prosthesis	Amelogenin (184 bp in D4S43 locus)	Resin prosthesis used in oral cavity and left at room temperature for 200 days can be used for DNA extraction and analysis.

(Continued)

Table 3. Continued

Studied by	Study Samples, Design and Description	DNA Primer Used	Outcome
Murakami et al. (2000) [49]	Freshly extracted primary (n = 6) and permanent teeth (n = 16); pulp stored in seawater (n = 8); Teeth stored in soil 8 teeth for 1 week ;8 teeth for 4 weeks and 6 teeth for 8 weeks along with 3 cases	Alphoid repeat primers	100% result; 1/6 immersed teeth in sea water for 4 weeks, successful gender determined through DNA hard tissue.
Sivagami et al. (2000) [68]	10 known (6 males and 4 females) and 10 unknown samples	Amelogenin	Despite Ultrasonic treatment of teeth samples, sufficient yield amount with good quality DNA obtained. AMEL gene proves to be good primer for gender determination in Indian population.
Tsuchimochi et al. (2002) [69]	Gender determination from human dental pulp incinerated to various temperature (200, 300, 400 and 500 degrees Celsius)	Alphoid repeat primers, STR and Y loci (DYZ3, DYS19, DYS390, DYS393)	All markers typed in samples incinerated upto 300°C for 2 min.
Andreasson et al. (2003) [70]	DNA extraction from teeth using Real time DNA Quantification Assay; STR analysis	Amelogenin	
Bilge et al. (2003) [71]	Case report study of age and gender through DNA profiling	Amelogenin	Male gender identification could be done successfully.
Chang et al. (2003) [72]	Gender analysis in 338 male individuals in a Malaysian population comprising Malays, Chinese and Indians, using the AmpFISTR Profiler Plus kit	Amelogenin	The amelogenin test gave a significant proportion of null alleles in the Indian ethnic group (3.6% frequency) and 0.88% frequency in the Malay ethnic group due to a deletion of the gene on the Y chromosome. This sex test also failed in a forensic casework sample. Inclusion of amelogenin with MSY1 loci on short arm of Y chromosome is recommended for accurate result.
William et al. (2004) [73]	Gender typing from deciduous teeth exposed to incineration (100-500 degrees temperature)	Amelogenin	Successful identification from deciduous teeth incinerated at 200 degrees Celsius and below. Greater sensitivity using fragment gel analysis by laser induced fluorescence which achieved sex identification from same human teeth to 400 degrees Celsius.
Yoshida et al. (2005) [74]	80 fresh and 15 old teeth which were preserved less than 186 days through VNTR	Amelogenin (DYZ1, DYZ3 combined with DXZ1)	Successful sex determination from fluorescent probes and amelogenin even with low quantity DNA.
Kumar et al. (2005) [75]	Gender identification from Exfoliated primary teeth using phenol-chloroform method	Alphoid repeat primers (X and Y specific sequences, 131cbp and 172 bp)	Inferred that PCR analysis not effective after 6 months of extraction.
Anzai et al. (2005) [15]	Twenty saliva samples collected;5 of these samples were deposited on skin and then collected subjected to STR analysis	Multiple loci (A set of 15 STRs)	Amount od DNA recovered when deposited on skin was 14 times less than salivary DNA. Standardized protocol for DNA collection and extraction from skin-deposited saliva can be used as a method to recover salivary DNA in criminal cases.
Zhang et al. (2006) [76]	Archeological samples (n = 8 graves; total of 16 incinerated teeth)	Amelogenin	7/16 samples gave positive result; Ancient DNA from teeth are more marked than bone.
Kovatsi et al. (2009) [77]	Effectiveness of DNA repair protocol in improving genetic testing in ancient teeth sample (n = 10)	Amelogenin	Successful in 8/10 cases.
Gharib et al. (2010) [78]	Sex determination in tooth samples in Iraqi population (10 male; 10 female)	Amelogenin	Amelogenin proved as good marker for sex determination in Iraqi population.

(Continued)

Table 3. Continued

Studied by	Study Samples, Design and Description	DNA Primer Used	Outcome
Reddy et al. (2011) [17]	Determined gender from epithelial cells from tooth brush samples using real time PCR (n = 30)	SRY	100% sensitivity and 73.3% specificity; All male samples gave positive result while 4 out of 15 female sample gave false positive result.
Raimann et al. (2012) [79]	Gender determination from unidentified cadaveric samples (n = 26) samples included premolar and molar with different post-mortem interval	Y- STR	Successfully determined gender in all samples.
Vemuri et al. (2012) [80]	Extracted teeth subjected to various temperature (100, 200, 300 and 400 degree Celsius); seawater (20–36) and burying at 30 cm depth (n = 15)	Alphoid repeat primers	Out of 15 samples, 13 samples were successfully identified. No DNA yield at high temperature.
Khare et al. (2012) [81]	Saliva and blood sample from 20 subjects were collected	Amelogenin	Quantity of salivary DNA was less compared to blood DNA but sufficient for gender determination.
Laverde et al. (2013) [82]	Retrospective study of 300 forensic cases were devised (n = 220 males; 80 females)	Amelogenin	Male component in amelogenin marker may not always be successful in correct gender determination.
Zapico et al. (2013) [83]	Teeth samples from Dentin and pulp using silica based DNA extraction technique	Amelogenin	DNA yield depends on type of tooth (smaller teeth with smaller yield).
Zagga et al. (2014) [52]	Dry teeth samples from cadavers, primary and permanent teeth (n = 3 each)	Alphoid repeat	Successful use of alphoid repeat primers in dry teeth in Sokoto, Northwestern Nigeria population.
Marjanovic et al. (2015) [84]	Teeth samples from skeletal remains of World War II victims (n = 9); 12 buccal swabs from potential living relatives were collected.	PowerPlex Y STR kit(Including Amelogenin)	Profile success rate was 90%. Referent samples success rate was 100%.
Thangraj et al. (2016) [85]	A total of 40 samples were subjected for gender determination through molecular technique (20 male; 20 female)	Amelogenin	100% specificity and sensitivity.
George et al. (2016) [86]	Gender estimation from epithelial cells adhered to acrylic dentures (n = 38; 21 males and 17 females)	SRY	100% accuracy.
Datta et al. (2017) [87]	Teeth subjected to various environmental conditions (n = 50)	Amelogenin	
Kholief et al. (2017) [88]	Freshly extracted teeth were subjected for PCR analysis (n = 40)	Amelogenin	Ultrasonication of teeth samples yield sufficient amount of good quality. Amelogenin gene is good primer for sex determination in Iraqi population.
Muthuswamy et al. (2017) [53]	Maxillary and mandibular premolar (n = 10; 5 male and 5 females)	SRY	Prediction rate was 100% for male and 80% for female.
Chowdhury et al. 2018 [88]	130 extracted premolars put in different environmental conditions	Amelogenin	Steady decrease in the amount of DNA obtained on increasing the time period at which teeth were exposed, irrespective of the environmental condition. Successful gender determination by AMEL gene with complete sensitivity and specificity, except in temperatures of 350°C.
Bharath et al. 2019 [89]	30 partial dentures	SRY	Sex determination was possible from all samples with 100% accuracy.
Lim et al. 2019 [90]	17 teeth samples was subjected to burning at different temperatures ranging from 100°C to 500°C, at 2 to 10 minutes	Amelogenin	76.47% accuracy; 63.64% sensitivity.

chances of cross contamination and vulnerability to microbial action or decay.

The epithelial cells adhering to the toothbrush were also studied and proved as a promising source of DNA for sex determination [17]. Tanaka et al. demonstrated 10.430 $\mu\text{g}/\mu\text{l}$ DNA yield from toothbrushes [18]. Teeth are the best material for forensic studies due to their resistant nature against environmental assaults, such as; burn, dipping, trauma, damage, decay, and microbial degradation, teeth epitomize as an outstanding source of DNA [7].

Sweet et al. used cryogenic grinding of teeth to obtain DNA yield of 18.4 micrograms per tooth [19]. They exposed the identity of a homicide victim who was burned to death. DNA was retrieved from dental pulp (1.35 μg) extracted from intra osseous third molar. Potschet et al. obtained 6 μg –50 μg of genomic DNA from dental sample [7]. Prachi et al. retrieved 25 to 27 $\text{ng}/\mu\text{l}$ from 20 teeth samples [20]. With higher yield of DNA integrated modern analysis techniques gender can be identified from tooth samples with DNA from samples as little as 10 cells.

Amelogenin primer from bones and teeth has also been used to identify gender. Meyer et al. evaluated gender using amelogenin primer of 4000 and 7000 years old ancient bodies [21]. Similarly, Faerman et al. determined gender using amelogenin primer from bones and teeth in 200 to 8000 years old bodies [22]. After 9/11 attacks (2005) in about 20,000 human remain specimens, advanced STR-based technology was employed aiding in identification even scarce samples of bone and teeth [20].

DNA PROFILING SYSTEMS

Wyman and White (1980) pioneered the discovery of polymorphic locus in the human genome using a DNA probe and advanced techniques that evolved with time. DNA Profiling includes a range of typing systems. The molecular ways aiding forensic odontologist to determine gender from DNA are further discussed as under.

Polymerase Chain Reaction (PCR)

PCR was demonstrated first by Saiki et al. and its application in DNA synthesis by Mullis and Faloona [23, 24]. Currently, PCR has become indispensable tool in forensic especially for amplification of victim/suspect nucleic acid sequences and their subsequent matching in minimal sample cases. PCR is an in-vitro technique for multiplication of particular DNA sequences using enzymes by concurrent primer extension of the complementary strands of DNA. The use of PCR was limited until heat stable polymerases became available. Most PCR based techniques ease standardization, increase sensitivity of result and analyses old or badly degraded samples easily [25]. The major limitation of PCR technique is difficulty in DNA quantification [26]. This limitation was overcome by evolution of

REAL TIME PCR [27]. With advancements in reagents like the 'AmpFISTR MiniFiler' PCR amplification kit promising result even in case of degraded or adulterated profiling DNA samples becomes possible [28].

Southern blotting with single locus and multi locus DNA probes is the conventional technique, which is still in use in few research set-ups [29]. Initially multi-locus probes were employed for forensic genetic analysis but gradually these were substituted by VNTR using single locus polymorphisms under highly stringent conditions.

Recently, the advent of STR analysis has overpowered VNTR as method of choice for forensic identification.

Restriction Fragment Length Polymorphisms (RFLP)

This method employs "restriction endonuclease" enzyme, which cleaves DNA at specific site known as 'Palindromes' and then analyzes variable lengths of the DNA fragments thus obtained. The drawback with this typing system is that they require large amount of DNA samples, longer time and cannot be used with environmentally degraded samples. It is an older DNA typing systems and with the development of more efficient typing systems, it is no longer used [30].

Short Tandem Repeat (STR) Typing

STR is described as repetitive short sections of DNA at variable sites in the entire human genome. STR Typing intends to assess specific loci within nuclear DNA and owing to individual's variability in the number of repeats, STR analysis is considered as the most promising genetic typing method particularly for ancient samples [20, 31, 32]. Routinely, 9–16 sequences in a sample are assessed and D21S11, D7S820, TH01, D13S317, and D19S433 are common markers used in cases of doubtful parentage claims [33]. The accuracy rate increases with the length of sample.

Tierney and Bird integrated amelogenin assay into multiplex STR typing system facilitating simultaneous amplification of X and Y correlative amelogenin gene with various other STRs in single reaction leading to gender and STR genetic fingerprint of an individual to be concurrently acquired [34].

STR was firstly analyzed by electrophoretic system followed by fluorescent-based technology and use of DNA sequencers. This allows typing of large multiplexes (upto 10 systems) and automation of the technique [25]. Several commercial multiplexes are now available like SGM Plus (Applied Biosystems) having 10 loci including amelogenin, Promega (Madison, WI) multiplexes using manual electrophoretic systems or monochromatic sequencing platforms and the recent 15-plexes systems. The Poweplex16 (Promega) and the Identifier (Applied Biosystems) are the more frequently used systems [25].

Y-Chromosome STR Analysis

Y chromosome analysis can especially be used in paternity disputes since the Y chromosome passes directly from father to son. It can also be used in criminal cases when male suspect is involved in sexual assault cases. Y-STR evaluations are predominantly useful to identify the male DNA segment.

Joint Y chromosome (STR) plus MiniSTR technique was employed for identification from the DNA isolated from the teeth and bone remnants of World War II casualty's [35]. Y STR technique has been successfully commissioned in human skeletal remains identification unearthed from the corpus in Croatia, Bosnia and Herzegovina [36].

However, there are limitations of Y-chromosome analysis. Firstly, similar Y STR silhouette will be seen in all the male relatives for generations. Secondly, such strikingly similar profile is inherited exclusively patrilineally and that too in haploid fashion. The Y-PLEX 12, a commercial kit system, allowing co-amplification of eleven polymorphic STR loci, residing on the Y chromosome including amelogenin permits easier identification [37].

X-chromosome STR (X-STR) Analysis

X-STR are easy amplifiable, highly sensitive alleles and forms approved system particularly in doubtful parent-hood examination [38].

Single Nucleotide Polymorphism (SNP)

Single nucleotide polymorphisms (SNP) are the utmost prevalent type of genetic deviation between individuals. Each SNP represents different nucleotide in place of normal [39]. Small size, ability to analyze highly degraded samples, lower mutation rate are the reasons SNPs are gaining importance [40].

In recent years, SNP technique was effectively employed to identify 9/11 and Tsunami victims [41]. Pakstis et al have formulated world wide accepted 92 SNPs for individual identification [25]. Limitations of SNPs comprise, firstly, there are no widely established core loci, and, secondly, prerequisite of huge multiplexing assays for analysis of multiple SNPs simultaneously [42]. Though SNPs is the most promising and future of DNA technology, they are improbable to substitute STRs presently utilized in national DNA databases, Primarily, owing to small mutation rate, and, secondly, due to use of manifold typing programs making worldwide acceptable SNP assortment challenging [36]. Attempts are being made to explore the possibility to make SNP a substitute to STR [36].

GENES FOR GENDER DETERMINATION

DNA-based sex assessment can be executed by numerous sex typing genes and markers like zinc finger genes

Locus	Primers	Product Size
Amelogenin	5'-CCCTGGGCTCTGTAAAGAATAGT-3' 5'-ATCAGAGCTTAACTGGGAAGCTG-3'	X = 106 bp Y = 112 bp
Amelogenin	5'-ACCTCATCCTGGGCAACCCTGG-3' 5'-AGGCTTGAGGCCAACCATCAG-3'	X = 212 bp Y = 218 bp
Amelogenin	5'-CTGATGGTTGGCCTCAAGCCTGTG-3' 5'-TAAAGAGATTCAATTAACCTGACTG-3'	X = 977 bp Y = 788 bp
Amelogenin	5'-CCCTTTGAAGTGGTACCAAGCA-3' 5'-GCATGCCTAATATTTTTCAGGGAATA-3'	X = 80 bp Y = 83 bp
Centromeric alphoid repeat	5'-TATTTGGACTCTCTGTAGGA-3' (X3) 5'-TTCTACTACAAGGGTGTTCGA-3' (X4) 5'-GTGTAATTCACCTCCGGGAG-3' (Y3) 5'-ACAAAAGGTTCAATTCTGTGAG-3' (Y4)	X = 157 bp Y = 200 bp
Centromeric alphoid repeat	5'-AATCATCAAATGGAGATTG-3' (X1) 5'-GTTCAAGCTCTGTGAGTGAAA-3' (X2) 5'-ATGATAGAAACGGAAATATG-3' (Y11) 5'-AGTAGAATGCAAAAGGGCTC-3' (Y22)	X = 170 bp Y = 130 bp
ZFX/ZFY zinc finger gene	5'-CTGGAGAGCCCAAGCTGAC-3' 5'-TTGCTGTGACTGCCAAGAG-3' X/Y = 209bp after <i>Hae</i> III cuts	X = 172 +37 Y = 88 +84 +37
SRY	5'-ATAAGTATCGACCTCGTCGGAAG-3' 5'-GCACTTCGCTGCAGAGTACCAGG-3'	Y = 93 bp

Figure 1. Sequence for primers in amplifying amelogenin, centromeric alphoid repeats, ZFX/ZFY) and the SRY gene (Adapted from <https://strbase.nist.gov/sextype.html>).

ZFX/ZFY, SRY, centromeric alphoid repeats and the amelogenin gene [43]. Amelogenin is most widely used genetic marker for gender determination. Refer Figure 1.

Zinc Finger Genes

These sex-determining genes located on short arm of Y chromosome, codes for zinc finger protein. It comprises of 30 amino acid repeating region and acts as transcriptional activator proteins, which gets attached to particular DNA segment. The zinc-finger structure is reinforced by the zinc ion that synchronizes cysteine and histidine in diverse amalgamations. Crystallographic studies of conventional zinc-finger domains revealed presence of two β -sheets and one α -helix [44].

These are also referred as Testis determining factor (TDF) located in the genome of the heteromorphic individual and causes the bi-potential gonadal primordia to differentiate into testes. In addition a homologue of ZFY is present on the X chromosome and referred as ZFX, which codes for zinc finger protein and cross-hybridizes to ZFY under stringent conditions [45]. ZFY could be a useful marker for Y chromosome for gender screening [25, 46]. Pilly and Kramer successfully determined gender from pulp for 21 males and 24 females using ZFX and ZFY genes [47].

Centromeric Alphoid Repeats

In all human chromosomes, Alpha satellite DNA is positioned in the peri-centromeric region of all human chromosomes spaced at distance. It is comprised of a basic 171 base pairs long unit that is structured into tandemly arranged higher order repeats (HORs); HORs consists of variable amounts of basic repeats, starting with 4 (chromosome 2) and extending till 34 (chromosome Y) [48]. The HORs of a given chromosome are highly similar. The lengths of the alphoid repeats are variable. The alphoid repeat primers

are more accurate in sex estimation because both the X chromosome specific alphoid repeat sequence along with Y chromosome specific repeat sequence can be concurrently spotted [49].

Witt and Erickson for the first time used alphoid repeat primers for gender determination on dried human blood samples [50]. Hanaoka & Minagouchi also determined gender from alphoid repeat sequences from teeth [51]. Murakami et al. determined gender employing DNA extraction from freshly extracted permanent teeth pulp and dentin using X and Y chromosome-specific alphoid repeats sequences. These tooth were preserved at room temperature for about two decades. They also analyzed pulp of teeth that were immersed in seawater; the gender was successfully assessed in all eight teeth submerged for one week and in five out of six teeth submerged for four weeks. Similar interesting results were obtained for teeth stored in soil exhibiting positive gender assessment from all eight teeth buried for one week, seven out of eight teeth submerged for four weeks, and from all six teeth buried for eight weeks.

In another experiment, the gender of a mummified body could be determined even approximately one year after death only through dental pulp. Also sex determination was successfully achieved through analysis of dental hard tissue in the carcasses of 2 bodies kept under water for about one year and approximately eleven years wherein the pulp tissues had been dissolved and lost [42]. Zagga et al. studied molecular sex determination on dry teeth specimens from cadaveric, primary and permanent teeth samples using alphoid repeat primers and obtained 100% result [52].

SRY Genes

The *SRY* gene located on Yp11.2 from base pairs 2,786,855 to 2,787,741 encodes 'Sex-determining region Y' protein that plays role in sexual development of males [48]. This protein is a transcription factor and modulates the activity of specific genes to initiate the development that instigate a fetus to grow male gonads and avert the growth of female reproductive structures.

Muthuswamy determined gender from incinerated teeth using *SRY* gene [53]. However, certain syndromes such as maternal – fetal micro-chimerism, Turner syndrome, Klinefelter syndrome, Swyer syndrome or in partial gonadal dysgenesis can generate false positive result [48]. Nicole et al. also revealed that sex determination could be inaccurate in samples acquired from bone marrow transplanted subjects or in pregnant female carrying a male fetus [50, 54].

Amelogenin

Amel gene encodes amelogenin proteins, which comprises 90% of the total enamel matrix proteins and performs

a pivotal role in enamel morphogenesis and mineralization. The location of human amelogenin gene is on X and Y-chromosome at Xp22.1–p22.3 and Yp11.2 respectively [55]. On X chromosome it is 2872 base pairs long while on Y chromosome it has a size of 3272 base pairs. Almost 90% of the *amel* gene transcripts are articulated from the X-chromosome while the remaining from the Y-chromosome [56]. The X and Y copies of amelogenin gene do not undertake recombination that is correlative, and that is why, it is the most favored genetic marker for gender assessment in current scenario [52]. Nakahori et al. sequenced Amelogenin gene for the very first time [57]. Komuro T et al. used capillary gel electrophoresis method to determine gender through X and Y locus from DNA obtained from dental pulp tissue [58]. Many marketing kits that provide promising result are now accessible for gender assessment [59]. However, false positive result using amelogenin primer can be seen in cases of chimerism that is, bone marrow transplants or microchimerism such as pregnant women with male foetus, genetic miscellany cases, intersex conditions and trans-sexualism, or aneuploidy in the sexual chromosome [60].

Young et al. exposed an aberration with Klinefelter Syndrome in two undiagnosed males while using amelogenin locus and X and Y STR's. Michael and Brunner reported the first case of failure of an amelogenin gender test on a phenotypically normal male in the Israeli population [61]. Since single gene gave erroneous results, Yamamoto and Morikawa proposed a novel technique for gender assessment based on findings of *SRY*, *STS* and amelogenin gene regions with simultaneous amplification of their similar sequences by using multiplex PCR [62].

Table 3 highlights studies employing molecular techniques in samples from the oral cavity to determine gender.

CONCLUSION

Gender determination using molecular techniques in forensic odontology show prodigious outcome and hence has befitted as frontier in forensic odontology. Identification of an individual in mass disaster and crime investigation cases in current scenario is an ardent but highly desirable task. Thorough knowledge of forensic genetics is essential and is the need of the hour. Recent advancements in molecular technique for DNA typing useful in gender determination are both economical and time savers. These advancements will bring justice to people and large benefit to the society. However, still there are issues with technical expertise and dissemination of knowledge. Moreover, technologies like Next gen Sequencing and nanotechnology are bringing revolutions in human sciences but forensic odontology is yet to see their ride.

Declaration of interest: The Authors declare that there are no potential conflicts of interest.

Funding Source: None.

Acknowledgements: Department of Health and Research-Multidisciplinary Research Unit, King George's Medical University Lucknow, Uttar Pradesh, India

Authors Contribution: RP and SS have contributed in acquisition of data, analysis and interpretation of data, drafting of the article, critical revision and final approval of the submitted version. RD, EA, DM and SC have contributed in drafting of the article, critical revision and final approval of the submitted version.

REFERENCES

- [1] Kanchan, T., Krishan, K. (2013) Personal Identification in Forensic Examinations. *Anthropol* 2: 114. Doi: 10.4172/2332-0915.1000114
- [2] Sharma, S., David, M.P., Indira, A.P. (2017). Gender determination from dental pulp tissue. *International Journal of Contemporary Medical Research* 4(4):822-5.
- [3] da Silva, R.H., Sales-Peres, A., de Oliveira, R.N., de Oliveira, F.T., Sales-Peres, S.H. (2007). Use of DNA technology in forensic dentistry. *Journal of applied oral science: revista FOB*, 15(3), 156-61. <https://doi.org/10.1590/s1678-77572007000300002>
- [4] Calabrez, M.C.T., Saldanha, P.H. (1997) A pesquisa de DNA em odontologiaforense. In: Silva M. *Compêndio de odontologia legal*. Rio de Janeiro: Editora Medsi. p. 167-221.
- [5] Mayall, S.S., Agarwal, P., Vashisth, P. (2013) Dental DNA fingerprinting in identification of human remains. *Annals of Dental Specialty* 1(1):16-19.
- [6] Oliveira, R.N., Daruge, E., Galvão, L.C.C., Tumang, A.J. (1998) Contribuição da odontologia legal para identificação "post-mortem". *Rev Bras Odontol*. 55(2):117-22.
- [7] Manjunath, B.C., Chandrashekar, B.R., Mahesh, M., Vatchala Rani, R.M. (2011) DNA profiling and forensic dentistry—a review of the recent concepts and trends. *J Forensic Leg Med* 18(5):191-7. doi: 10.1016/j.jflm.2011.02.005.
- [8] Jeffreys, A.J., Brookfield, J.F., Semeonoff, R. (1985) Positive identification of an immigration test-case using human DNA fingerprints. *Nature* 317:818e9.
- [9] Lessig, R., Edelmann, J. (1995) Individualisation of dental tissue – an aid for odontological identification? *J Forensic Odontostomatol* 13:1-3.
- [10] Malaver, P.C., Yunis JJ (2003) Different dental tissues as source of DNA for human identification in forensic cases. *Croat Med J* 44(3):306-9.
- [11] Van-Wieren-de-Wijer, D.B., Maitland-van-der-Zee, A.H., de Boer, A., et al. (2009) Determinants of DNA yield and purity collected with buccal cell samples. *Eur J Epidemiol* 24:677-82.
- [12] Ghatak, S., Muthukumar, R.B., Nachimuthu, S.K. (2013) A simplified method for DNA extraction for human samples from PCR-RFLP Analysis. *Journal of Biomolecular Techniques* 24:224-31.
- [13] Walsh, D.J., Corey, A.C., Cotton, R.W. et al. (1992) Isolation of deoxyribonucleic acid (DNA) from saliva and forensic science samples containing saliva. *J Forensic Sci* 37:387-95.
- [14] Koh, D., Ng, P.K., Choo, S.G.L., Ng, V., Fu, Q. (2004) Effect of storage conditions on the extraction of PCR-quality genomic DNA from saliva. *Clin Chim Acta* 343: 191-4.
- [15] Anzai-Kanto, E., Hirata, M.H., Hirata, R.D.C., Nunes, F.D., Melani, R.F.H., Oliveira, R.N. (2005) DNA extraction from human saliva deposited on skin and its use in forensic identification procedures. *Braz Oral Res* 19(3):216-22.
- [16] George, R., Sriram, G., Saraswati, T.R., Sivapathasundaram, B. (2010) Isolation of epithelial cells from acrylic removable dentures and gender identification by amplification of SRY gene using real time PCR. *J Forensic Dent Sci* 2(1):32-6.
- [17] Reddy, V.S.A., Sriram, G., Saraswati, T., Sivapathasundaram, B. (2011) Isolation of epithelial cells from tooth brush and gender identification by amplification of SRY gene. *J Forensic Dent Sci* 3(1):27-32.
- [18] Tanaka, M., Yoshimoto, T., Nozawa, H., Ohtaki, H., Kato, Y., et al. (2000) Usefulness of a toothbrush as a source of evidential DNA for typing. *J Forensic Sci* 45(3): 674-6.
- [19] Sweet, D., Sweet, C.H.W. (1995) DNA analysis of dental pulp to link incinerated remains of homicide victim to crime scene. *J Forensic Sci* 40:310-4.
- [20] Naik, P.R., Acath, D.D., Sharma, G.H., Neelkarv, A.R. (2012) Viability of human dental pulp in determination of sex of an individual by identifying by SRY gene through DNA analysis: A single blind pilot study. 24(2):133-6.
- [21] Meyer, E., Wiese, M., Bruchhaus, H., Claussen, M., Klein, A. (2000) Extraction and amplification of authentic DNA from ancient human remains. *Forensic Sci Int*.113: 87-90.
- [22] Faerman, M., Filon, D., Kahila, G., Greenblatt, C.L., Smith, P., Oppenheim, A. (1995) Sex identification of archaeological human remains based on amplification of the X and Y amelogenin alleles. *Gene*.167: 327-32.
- [23] Saiki, R.K., Scharf, S., Faloona, F. et al. (1985) Enzymatic amplification of ss-globin sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* 230:1350-4.
- [24] Mullis, K.B., Faloona, F.A. (1987) Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. *Methods Enzymol* 155:335-50.
- [25] Pakstis, A.J., Speed, W.C., Fang, R., et al. (2010) SNPs for a universal individual identification panel. *Hum Genet* 127:315e24.
- [26] Kubista, M., Andrade, J.M., Bengtsson, M., Forootan, A., Jonák, J., Lind, K., et al. The real-time polymerase chain reaction. *Mol Aspects Med* 2006;27:95e125.
- [27] Higuchi, R., Fockler, C., Dollinger, G., Watson, R. (1993) Kinetic PCR analysis: real-time monitoring of DNA amplification reactions. *Biotechnol (NY)* 11:1026e30.
- [28] Alenizi, M.A., Goodwin, W., Hadi, S., Alenizi, H.H., Altamar, K.A., Alsik, M.S. (2009) Concordance between the Amp FISTR Mini Filer and Amp FISTR identifier PCR amplification kits in the Kuwaiti population. *J Forensic Sci* 54:350e2.
- [29] Carracedo, A., Sanchez-Diz, P. (2005) Forensic DNA typing Technologies-A review. *Methods and Molecular Biology*. 1-278.
- [30] Zehner, R., Zimmermann, S., Mebs, D. (1998) RFLP and sequence analysis of the cytochrome b gene of selected animals and man: Methodology and forensic application. *Int J Leg Med* 111:323-7.
- [31] Bramanti, B., Hummel, S., Chiarelli, B., Herrmann, B. (2003) Ancient DNA Analysis of the Delta F508 Mutation, *Human Biology* 75(1): 105-115.
- [32] Butler, J. M. (2005) *Forensic DNA Typing: Biology, Technology and genetics of STR markers*, 2nd edition, Elsevier Academic press, New York.
- [33] Jayanandan, M., Gopalkrishnan, S. (2011) Practical aspects of DNA based study in forensic dentistry. *JFDS* 3(1): 38-45.
- [34] Tierney, S., Bird, J. (2014) Sex Identification Of Human Remains From An Irish Medieval Population Using Biomolecular methods. *European Scientific Journal* 2:1857-81.
- [35] Definis, G.M., Sutlovic, D. (2007) Skeletal remains from World War II mass grave: from discovery to identification. *Croat Med J* 48:520e7.
- [36] Andelinovic, S., Sutlovic, D., ErcegIvkovic, I., et al. (2005) Twelve-year experience in identification of skeletal remains from mass graves. *Croat Med J* 46:530e9.
- [37] Shewale, J.G., Nasir, H., Schneida, E., Gross, A.M., Budowle, B., Sinha, S.K. (2004) Y-chromosome STR system, Y-PLEX 12, for forensic casework: development and validation. *J Forensic Sci* 49:1278e90.
- [38] Datta, P., Datta, S.S. (2012) Role of deoxyribonucleic acid technology in forensic dentistry. *J Forensic Dent Sci*. 4(1): 42-6. doi: 10.4103/0975-1475.99165.
- [39] Genetic Home references. (2018) United States National Library of Medicine. available on: https://ghr.nlm.nih.gov/primer/genomic_research/snp
- [40] Kim, J.J., Han, B.G., Lee, H.I., Yoo, H.W., Lee, J.K. (2010) Development of SNP-based human identification system. *Int J Leg Med* 124:125e31.
- [41] Gill, J.R. (2006) 9/11 and the New York City Office of Chief medical Examiner. *Forensic Sci Med Pathol* 2:29e32.

- [42] Butler, J.M., Budowle, B., Gill, P. et al. (2008) Report on ISFG SNP panel Discussion. *Forensic Sci Int (suppl 1)*:471 e2.
- [43] Muthuswamy, D. (2017) Determination of Sex of an Individual by Identifying SRY Gene through PCR Analysis of DNA Extracted From Incinerated Teeth-A Single Blind Pilot Study *Int J foren Sci* 2(3):000128.
- [44] Cassandri, M. (2017) Zinc-finger proteins in health and disease *Cell Death Discov* 13;3:17071.
- [45] Aasen, E., Medrano, J. F. (1990) Amplification for ZFY and ZFX genes for sex identification in humans, cattles, sheep and goats. *Biotechnology, Nature Publishing Group* 8:1279–81.
- [46] Shaw, C.N., Wilson, P.J., White, B.N. (2003) A Molecular method for gender determination in Mammals. *Journal of Mammalogy* 84(1):123–8.
- [47] Pilly, U., Kramer, B. (1997) A simple method for the determination of sex from the pulp of freshly extracted human teeth utilizing the polymerase chain reaction. *J Dent Assoc S Afr* 52(11): 673–7.
- [48] Roizès, G. (2006) Human centromeric alphoid domains are periodically homogenized so that they vary substantially between homologues. Mechanism and implications for centromere functioning. *Nucleic Acids Res* 34:1912–24.
- [49] Murakami, H., Yamamoto, Y., Yoshitome, K., et al. (2000) Forensic study of sex determination using PCR on teeth samples. *Acta Med Okayama* 54:21–32.
- [50] Witt, M., Erickson, R.P. (1989) A rapid method for detection of Y-chromosomal DNA from dried blood specimens by the polymerase chain reaction. *Hum Genet* 82:271–4.
- [51] Hanaoka, Y., Minaguchi, K. (1996) Sex determination from blood and teeth by PCR amplification of the alphoid satellite family. *J Forensic Sci* 41(5):855–8.
- [52] Zagga, A., Ahmed, H.O., Ismail, S., Tadros, A. (2014) Molecular sex identification of dry human teeth specimens from Sokoto, North-western Nigeria. *J Forensic Dent Sci.* 6(2):132–8. doi: 10.4103/0975-1475.132544.
- [53] Muthusamy, D. (2017) Determination of Sex of an Individual by Identifying SRY Gene through PCR Analysis of DNA Extracted From Incinerated Teeth-A Single Blind Pilot Study. *Int J Foren Sci* 2(3):000128.
- [54] Costa, J.M., Benachi, A., Gautier, E., et al. (2001) First-trimester fetal sex determination in maternal serum using real-time PCR. *Prenat Diagn* 21(12):1070–1074.
- [55] Yamada, Y., Yamamoto, K., Yoshii, T., Ishiyama, I. (1989) Analysis of DNA from tooth and application to forensic dental medicine. *Nihon Hoigaku Zasshi*. 43(5):420–3.
- [56] Taylor, A., Filderman, A., Blumenfeld, A., et al. (2006) High yield of biologically active recombinant human amelogenin using the baculovirus expression system. *Protein Expr Purif* 45:43–53.
- [57] Nakahori, Y., Takenaka, O., Nakagome, Y. (1991) A human X-Y homologous region encodes amelogenin. *Genomics* 9:264–9.
- [58] Komuro, T., Nakamura, M., Tsutsumi, H., Mukoyama, R. (1998) Gender determination from dental pulp by using capillary gel electrophoresis of amelogeninlocus. *J Forensic Odontostomatol* 1998;16:23e6.
- [59] Pouchkarev, V.P., Shved, E.F., Novikov, P.I. (1998) Sex determination of forensic samples by polymerase chain reaction of the amelogenin gene and analysis by capillary electrophoresis with polymer matrix. *Electrophoresis* 19:76e9.
- [60] Von Wurmb-Schwark, N., Bosniki, H., Ritz-Timme, S. (2007) What do the X and Y chromosome tell us about sex and gender in forensic case analysis? *J. Forensic Leg Med* 14:27–30.
- [61] Pawlowski, R. (1995) Identification analysis using PCR of fresh or fixed and paraffin wax embedded tissue. *Pol Tyg Lek* 50(36–39): 16–20.
- [62] Yamamoto, K. (1996) Molecular biological studies on teeth, and inquests. *Forensic Sci Int.* 28;80(1–2):79–87.
- [63] Chen, L., Sun, G., Wu, M. (1994) Influence exerted by environmental and physicochemical factors on the results of sex identification of human dental pulp by polymerase chain reaction *Hua Xi Yi Ke Da Xue Xue Bao* 25(3):253–8.
- [64] Jiang, B., Guo, J.Y. (1997) Determination of sex using PCR of the X-Y homologous gene amelogenin *Fa Yi XueZaZhi* 13(2): 68–70, 128.
- [65] Ovchinnikov, I.V., Ovtchinnikova, O.I., Druzina, E.B., Buzhilova, A.P., Makarov, N.A. (1998) Molecular genetic sex determination of Medieval human remains from north Russia: comparison with archaeological and anthropological criteria. *Anthropol Anz.* 56(1): 7–15.
- [66] Urbani, C., Lastrucci, R.D., Kramer, B. (1999) The effect of temperature on sex determination using DNA-PCR analysis of dental pulp. *J Forensic Odontostomatol.* 17(2):35–9.
- [67] Inoue, M., Hanaoka, Y., Minaguchi, K. (2000) Personal identification by DNA analysis of samples from dental prostheses made of acrylic resin. *Bull Tokyo Dent Coll.* 41:175–85.
- [68] Sivagami, A.V., Rao, R. A., Varshney, U. (2000) A simple and cost-effective method for preparing DNA from the hard tooth tissue, and its use in polymerase chain reaction amplification of amelogenin gene segment for sex determination in an Indian population. *Forensic Science International* 110:107–115.
- [69] Tsuchimochi, T., Iwasa, M., Maeno, Y. et al (2002) Chelating resin-based extraction of DNA from dental pulp and sex determination from incinerated teeth with Y-chromosomal alphoid repeat and short tandem repeats. *Am J Forensic Med Pathol* 23(3):268–71.
- [70] Andréasson, H., Allen, M. (2003) Rapid quantification and sex determination of forensic evidence materials. *J Forensic Sci* 48(6): 1280–7.
- [71] Bilge, Y., Kedici, P.S., Alakoc, Y.D., Iku'er, K.Ü., Ilkyaz, Y.Y. (2003) The identification of a dismembered human body: a multidisciplinary approach. *Forensic Sci Int* 137:141–6.
- [72] Chang, Y.M., Burgoyne, L., Both, K. (2003) Higher Failures of Amelogenin Sex Test in an Indian Population Group. *Journal of Forensic Sciences* 48(6):1309–13.
- [73] Williams, D., Lewis, M., Franzen, T. et al. (2004) Sex determination by PCR analysis of DNA extracted from incinerated, deciduous teeth. *Sci Justice.* 44(2):89–94.
- [74] Yoshida, M., Suno, M., Asari, M. et al (2005) Sex identification from DNA in old teeth. *Hokkaido Igaku Zasshi* 80(2):191–9.
- [75] Kumar, M.G., Hegde, A.M. (2005) Sex identification from exfoliated primary teeth—a PCR study. *J Clin Pediatr Dent* 30(1):19–21.
- [76] Zhang, H.Q., Yang, Z.Q., Liu, F.E., Zhang, J., Zhao, W.M. (2006) Homologous amelogenin gene test of archaeological samples *Fa Yi Xue Za Zhi* 22(3):213–6.
- [77] Kovatsi, L., Nikou, D., Triantaphyllou, S., Njau, S.N., Voutsaki, S., Kouidou, S. (2009) DNA repair enables sex identification in genetic material from human teeth. *Hippokratia* 13(3):165–8.
- [78] Gharib, A.M. (2010) Use of teeth samples for gender determination by PCR in Iraqi case. *Iraqi J Biotech* 9:819–27.
- [79] Raimann, P.E., Picanço, J.B., d Silva, S.B.S., Albuquerque, T.C.K., Paludo, F.J.O., Alho, C.S. (2012) Procedures to recover DNA from premolar and molar teeth of decomposed cadavers with different post-mortem intervals. *Archives of Oral Biology* 57:1459–1466.
- [80] Vemuri, S., Ramya, R., Rajkumar, K., Rajashree, P. (2012) Influence of various environmental conditions on DNA isolation from dental pulp for sex determination using Polymerase Chain Reaction. *SRM Journal of Research and Dental Sciences* 3(4):231–235.
- [81] Khare, P., Raj, V., Chandra, S., Agarwal, S. (2014) Quantitative and qualitative assessment of DNA extracted from saliva for its use in forensic identification. *J Forensic Dent Sci* 6:81–5.
- [82] Laverde, L. (2013) Sex determination problem in forensic genetic analysis. *Forensic Science International. Genetics Supplement* 4:e350–e351.
- [83] Zapico, S.C., Ublekar, H. (2013) Sex determination from dentin and pulp in a medico legal context. *JADA* 144(12):1379–1385.
- [84] Marjanovic, D., Metjahic, N.H., Cakar, J. et al. (2015) Identification of human remains from the Second World War mass graves uncovered in Bosnia and Herzegovina. *Croat Med J.*;56:257–62.
- [85] Praveen, S.T., Aswath, N. (2016) DNA isolation from teeth by organic extraction and identification of sex of the individual by analyzing the AMEL gene marker using PCR. *J Forensic Dent Sci* 8:18–21.

- [86] Datta, P., Bhosale, S., Singh, R. et al. (2017) Amelogenin: The pioneer in Gender determination from Forensic Dental Samples . *J Clin Diagn Res* 11(2);56–9.
- [87] Kholief, M., Shanawany, S.E., Gomma, R. (2017) Sex determination from dental pulp DNA among Egyptians. *Egyptian Journal of Forensic Sciences* 7:29.
- [88] Chowdhury, R., Singhvi, A., Bagul, N., Bhatia, S., Singh, G., Goswami, S. (2018) Sex determination by amplification of amelogenin gene from dental pulp tissue by polymerase chain reaction. *Indian J Dent Res* 29(4):470–6. https://doi.org/10.4103/ijdr.IJDR_274_17
- [89] Bharath, T.S., Saraswathi, T.R., Suresh Sajjan, M.C., Ramchandran, C.R., Govindraj Kumar, N. (2019) Isolation and quantification of DNA from epithelial cells obtained from acrylic removable partial denture for sex identification. *Egypt J Forensic Sci* 9(1):19. <https://doi.org/10.1186/s41935-019-0123-9>
- [90] Lim, J.J.-Y., Khamis, M.F., Abd Rashid, N.H. (2019) Identification of amelogenin gene on burnt teeth samples through nested polymerase chain reaction amplification for sex identification. *Jurnal Sains Kesihatan Malaysia* 17(01):91–8. <https://doi.org/10.17576/jskm-2019-1701-12>